



Kinetic models for socio-economic dynamics of speculative markets

Dario Maldarella*, Lorenzo Pareschi

Department of Mathematics and CMCS, University of Ferrara, Via Machiavelli 35, I-44100 Ferrara, Italy

ARTICLE INFO

Article history:

Received 28 September 2010

Received in revised form 6 July 2011

Available online 7 September 2011

Keywords:

Kinetic models

Opinion formation

Stock market

Power laws

Behavioral finance

ABSTRACT

In this paper we introduce a simple model for a financial market characterized by a single stock or good and an interplay between two different trader populations, chartists and fundamentalists, which determine the price dynamics of the stock. The model has been inspired by the microscopic Lux–Marchesi model (Lux and Marchesi (2000, 1999) [3,25]). The introduction of kinetic equations permits to study the asymptotic behavior of the investments and the price distributions and to characterize the regimes of lognormal behavior and the formation of power law tails.

© 2011 Elsevier B.V. All rights reserved.

1. Introduction

Most speculative markets at national and international level share a number of stylized facts, like volatility clustering and fat tails of returns, for which a satisfactory explanation is still lacking in standard theories of financial markets [1]. Such stylized facts are now almost universally accepted among economists and physicists and it is now clear that financial market dynamics give rise to some kind of universal scaling laws.

Showing similarities with scaling laws for other systems with many interacting particles, a description of financial markets as multi-agent interacting systems appeared to be a natural consequence [2–6]. This topic was pursued by quite a number of contributions appearing in both the physics and economics literature in recent years [7–13,4,14,6]. This new research field borrows several methods and tools from classical statistical mechanics, where emerging complex behavior arises from relatively simple rules due to the interaction of a large number of components [15].

Starting from the microscopic dynamics, kinetic models can be derived with the tools of classical kinetic theory of fluids [7,16,17,10,18,12,19–24]. In contrast with microscopic dynamics, where behavior often can be studied only empirically through computer simulations, kinetic models based on PDEs allow us to derive analytically general information on the model and its asymptotic behavior.

In this paper we introduce a simple Boltzmann-like model for a speculative market characterized by a single stock and a socio-economical interplay between two different types of traders, chartists and fundamentalists. The model is strictly related to the microscopic Lux–Marchesi model [3,25] and to kinetic models of opinion formation recently introduced in Ref. [26]. In addition, we take into account some psychological and behavioral components of the agents, like the way they interact with each other and perceive the risk, which may produce nonrational behaviors. This is done by means of a suitable “value function” in agreement with the Prospect Theory by Kahneman and Tversky [27,28]. As we will show people systematically overreacting produces substantial instabilities in the stock market.

In an earlier paper [16] a similar approach has been used considering a single population of investors interacting in the stock market on the basis of the microscopic Levy–Levy–Solomon model [13]. The emergence of a lognormal behavior for the

* Corresponding author.

E-mail addresses: dario.maldarella@unife.it (D. Maldarella), lorenzo.pareschi@unife.it (L. Pareschi).

wealth distribution of the agents has been shown. Though the theoretical set-up of the analysis is close in certain respects to that of [16], the structure of the model is rather different. Namely, the description of individual behavior follows an opinion formation dynamics strictly connected with the price trend. In this way, the heterogeneity among agents as well as their social interactions will be taken into account both of which are key elements affecting the outcome of the overall market dynamics.

Following the analysis developed in Refs. [17,26], we shall prove that the Boltzmann model converges in a suitable asymptotic limit toward convection–diffusion equations of Fokker–Planck type. Other Fokker–Planck equations were obtained using different approaches in Refs. [7,29,30]. This permits to study the asymptotic behavior of the investments and the price distributions and to characterize the regimes of lognormal behavior and the ones with power law tails. The main finding of the present paper is that the presence of heterogeneous strategies, both fundamentalists and chartists, is essential to achieve basic stylized facts like the presence of fat tails.

The rest of the paper is organized as follows. In Section 2 we introduce the Boltzmann kinetic model for the interacting chartists and the price evolution. Details of the strategy exchange between chartists and fundamentalists are also presented here. A characterization of the admissible equilibrium states of the resulting system is then reported. Next, in Section 3, with the aim to study the asymptotic behavior of the chartists and price distributions, we introduce simpler Fokker–Planck approximations of the Boltzmann system and give explicit expressions of the long time behavior. The mathematical details of the derivation of such Fokker–Planck models are reported in separate appendices at the end of the manuscript. Numerical results which confirm the theoretical analysis are given in Section 4 and some concluding remarks are discussed in the last section.

2. A kinetic model for multiple agents interactions

We describe a simple financial market characterized by a single stock or good and an interplay between two different trader populations, chartists and fundamentalists, which determine the price dynamics of such stock (good). The aim is to introduce a kinetic description both for the behavior of the microscopic agents and for the price, and then to exploit the tools given by kinetic theory to get more insight about the way the microscopic dynamics of each trading agent can influence the evolution of the price, and be responsible of the appearance of ‘stylized’ facts like ‘fat tails’ and ‘lognormal’ behavior.

2.1. Kinetic setting

Similarly to Lux and Marchesi model [3,25], the starting point is a population of two different kind of traders, chartists and fundamentalists. Chartists are characterized by their number density ρ_C and the investment propensity (or opinion index) y of a single agent whereas fundamentalists appear only through their number density ρ_F . The value $\rho = \rho_F + \rho_C$ is invariant in time so that the total number of agents remains constant. In the sequel we will assume for simplicity $\rho = 1$.

Dynamics of investment propensity among chartists Let us define $f(y, t)$, $y \in [-1, 1]$, the distribution function of chartists with investment propensity y at time t . Positive values of y represent buyers, negative values characterize sellers and close to $y = 0$ we have undecided agents. Clearly

$$\rho_C(t) = \int_{-1}^1 f(y, t) dy.$$

Moreover we define the mean investment propensity

$$Y(t) = \frac{1}{\rho_C(t)} \int_{-1}^1 f(y, t) y dy. \quad (1)$$

For a given price $S(t)$ and price derivative $\dot{S}(t) = dS(t)/dt$ the microscopic dynamics of the investment propensity of chartists is characterized by the following binary interactions $(y, y_*) \rightarrow (y', y'_*)$ with

$$\begin{aligned} y' &= (1 - \alpha_1 H(y) - \alpha_2) y + \alpha_1 H(y) y_* + \alpha_2 \Phi \left(\frac{\dot{S}(t)}{S(t)} \right) + D(y) \eta, \\ y'_* &= (1 - \alpha_1 H(y_*) - \alpha_2) y_* + \alpha_1 H(y_*) y + \alpha_2 \Phi \left(\frac{\dot{S}(t)}{S(t)} \right) + D(y_*) \eta_*. \end{aligned} \quad (2)$$

Here $\alpha_1 \in [0, 1]$ and $\alpha_2 \in [0, 1]$, with $\alpha_1 + \alpha_2 \leq 1$, measure the importance the individuals place on other’s opinions and actual price trend in forming expectations about future price changes. The random variables η and η_* are assumed distributed accordingly to $\Theta(\eta)$ with zero mean and variance σ^2 and measure individual deviations from the average behavior. The function $H(y) \in [0, 1]$ is taken concave and symmetric on the interval I , and characterize the herding behavior, whereas $D(y) \geq 0$ defines the diffusive behavior, and will be also taken concave and symmetric on I . Simple examples of herding function and diffusion function are given by

$$H(y) = a + b(1 - |y|), \quad D(y) = (1 - y^2)^\gamma,$$

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات