

The power-law tail exponent of income distributions

F. Clementi^{a,b,*}, T. Di Matteo^b, M. Gallegati^c

^aDepartment of Public Economics, University of Rome “La Sapienza”, Via del Castro Laurenziano 9, 00161 Rome, Italy

^bApplied Mathematics, Research School of Physical Sciences and Engineering, The Australian National University, 0200 Canberra, Australia

^cDepartment of Economics, Università Politecnica delle Marche, Piazzale Martelli 8, 60121 Ancona, Italy

Available online 12 May 2006

Abstract

In this paper we tackle the problem of estimating the power-law tail exponent of income distributions by using the Hill’s estimator. A subsample semi-parametric bootstrap procedure minimizing the mean squared error is used to choose the power-law cutoff value optimally. This technique is applied to personal income data for Australia and Italy.

© 2006 Elsevier B.V. All rights reserved.

Keywords: Personal income; Pareto’s index; Hill’s estimator; Bootstrap

1. Introduction

Since Pareto it has been recognized that a *power-law* provides a good fit for the distribution of high incomes [1]. The Pareto’s law asserts that the complementary cumulative distribution $P_{>}(y) = 1 - \int_{-\infty}^y p(\xi) d\xi \rightarrow P_{>}(u)(u/y)^\alpha$, with $y \geq u$, where $u > 0$ is the threshold value of the distribution and $\alpha > 0$ turns out to be some kind of index of inequality of distribution. The fit of such distribution is usually performed by judging the degree of linearity in a double logarithmic plot involving the empirical and theoretical distribution functions, in such a way that the estimation of u of the distribution does not seem to follow a neutral procedure. Moreover, recent studies have criticized the reliability of this geometrical method by showing that linear-fit based methods for estimating the power-law exponent tend to provide biased estimates, while the maximum likelihood estimation method produces more accurate and robust estimates [2,3]. Hill proposed a conditional maximum likelihood estimator for α based on the k largest order statistics for non-negative data with a Pareto’s tail [4]. That is, if $y_{[n]} \geq y_{[n-1]} \geq \dots \geq y_{[n-k]} \geq \dots \geq y_{[1]}$, with $y_{[i]}$ denoting the i th order statistic, are the sample elements put in descending order, then the Hill’s estimator is

$$\hat{\alpha}_n(k) = \left[\frac{1}{k} \sum_{i=1}^k (\log y_{n-i+1} - \log y_{n-k}) \right]^{-1}, \quad (1)$$

*Corresponding author. Department of Public Economics, University of Rome “La Sapienza”, Via del Castro Laurenziano 9, 00161 Rome, Italy. Tel.: +39 06 49766843; fax: +39 06 4461964.

E-mail address: fabio.clementi@uniroma1.it (F. Clementi).

where n is the sample size and k an integer value in $[1, n]$. Unfortunately, the finite-sample properties of the estimator (Eq. (1)) depend crucially on the choice of k : increasing k reduces the variance because more data are used, but it increases the bias because the power-law is assumed to hold only in the extreme tail.

Over the last 20 years, estimation of the Pareto's index has received considerable attention in extreme value statistics [5]. All of the proposed estimators, including the Hill's estimator, are based on the assumption that the number of observations in the upper tail to be included, k , is known. In practice, k is unknown; therefore, the first task is to identify which values are really extreme values. Tools from exploratory data analysis, as the quantile-quantile plot and/or the mean excess plot, might prove helpful in detecting graphically the quantile $y_{[n-k]}$ above which the Pareto's relationship is valid; however, they do not propose any formal computable method and, imposing an arbitrary threshold, they only give very rough estimates of the range of extreme values.

Given the bias-variance *trade-off* for the Hill's estimator, a general and formal approach in determining the best k value is the minimization of the *Mean Squared Error (MSE)* between $\hat{\alpha}_n(k)$ and the theoretical value α . Unfortunately, in empirical studies of data the theoretical value of α is not known. Therefore, an attempt to find an approximation to the sampling distribution of the Hill's estimator is required. To this end, a number of innovative techniques in the statistical analysis of extreme values proposes to adopt the powerful bootstrap tool to find the optimal number of order statistics adaptively [6–9]. By capitalizing on these recent advances in the extreme value statistics literature, in this paper we adopt a subsample semi-parametric bootstrap algorithm in order to make a reasonable and more automated selection of the extreme quantiles useful for studying the upper tail of income distributions and to end up at less ambiguous estimates of α . This methodology is described in Section 2 and its application to Australian and Italian income data [10,11] is given in Section 3. Some conclusive remarks are reported in Section 4.

2. Estimation technique for threshold selection

In this section, we consider the problem of finding the optimal threshold u_n^* —or equivalently the optimal number k^* of extreme sample values above that threshold—to be used for estimation of α . In order to achieve this task, we minimize the *MSE* of the Hill's estimator (Eq. (1)) for a series of thresholds $u_n = y_{[n-k]}$, and pick the u_n value at which the *MSE* attains its minimum as u_n^* . Given that different threshold series choices define different sets of possible observations to be included in the upper tail of a specific observed sample $\mathbf{y}_n = \{y_i; i = 1, 2, \dots, n\}$, only the observations exceeding a certain threshold that are additionally distributed according to a Pareto's cumulative distribution function $PD_{\hat{\alpha}_n(k), u_n}(y)$ are included in the series. In order to check this condition, we perform for each threshold in the original sample a *Kolmogorov–Smirnov (K–S)* goodness-of-fit test for the null hypothesis $H_0 : \hat{F}_n(y) = PD_{\hat{\alpha}_n(k), u_n}(y)$ versus the general alternative of the form $H_1 : \hat{F}_n(y) \neq PD_{\hat{\alpha}_n(k), u_n}(y)$, where $\hat{F}_n(y)$ is the empirical distribution function, and $\hat{\alpha}_n(k)$ is a prior estimate for each threshold u_n of the Pareto's tail index obtained through the Hill's statistic. Following the methodology in [12], the formal steps in making a test of H_0 are as follows:

(a) Calculate the original *K–S* test statistic D by using the formula

$$D = \sup_{-\infty < y < \infty} |\hat{F}_n(y) - PD_{\hat{\alpha}_n(k), u_n}(y)|.$$

(b) Calculate the modified form T^* by using the formula

$$T^* = D \left(\sqrt{n} + 0.12 + \frac{0.11}{\sqrt{n}} \right). \quad (2)$$

(c) Reject H_0 if T^* exceeds the cutoff level, z , for the chosen significance level.

To obtain an estimate of finite-sample bias and variance (and thus *MSE*) at each threshold coming from the null hypothesis H_0 , a natural criterion is to use the *bootstrap* [13]. In its purest form, the bootstrap involves

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات