



Interactive fuzzy random cooperative two-level linear programming through level sets based probability maximization

Masatoshi Sakawa*, Takeshi Matsui

Faculty of Engineering, Hiroshima University, Higashi-Hiroshima 739-8527, Japan

ARTICLE INFO

Keywords:

Two-level linear programming problems
Fuzzy random variables
Level sets
Probability maximization
Interactive decision making

ABSTRACT

In this paper, assuming cooperative behavior of the decision makers, two-level linear programming problems under fuzzy random environments are considered. To deal with the formulated fuzzy random two-level linear programming problems, α -level sets of fuzzy random variables are introduced and an α -stochastic two-level linear programming problem is defined for guaranteeing the degree of realization of the problem. Taking into account vagueness of judgments of decision makers, fuzzy goals are introduced and the α -stochastic two-level linear programming problem is transformed into the problem to maximize the satisfaction degree for each fuzzy goal. Through probability maximization, the transformed stochastic two-level programming problem can be reduced to a deterministic one. Interactive fuzzy programming to derive a satisfactory solution for the decision maker at the upper level in consideration of the cooperative relation between decision makers is presented. An illustrative numerical example is provided to demonstrate the feasibility and efficiency of the proposed method.

© 2012 Elsevier Ltd. All rights reserved.

1. Introduction

In the real world, we often encounter situations where there are two or more decision makers in an organization with a hierarchical structure, and they make decisions in turn or at the same time so as to optimize their objective functions. In particular, consider a case where there are two decision makers; one of the decision makers first makes a decision, and then the other who knows the decision of the opponent makes a decision. Such a situation is formulated as a two-level programming problem (Sakawa & Nishizaki, 2009). In the context of two-level programming, the decision maker at the upper level first specifies a strategy, and then the decision maker at the lower level specifies a strategy so as to optimize the objective with full knowledge of the action of the decision maker at the upper level. In conventional multi-level mathematical programming models employing the solution concept of Stackelberg equilibrium, it is assumed that there is no communication among decision makers, or they do not make any binding agreement even if there exists such communication (Bialas & Karwan, 1984; Nishizaki & Sakawa, 2000; Shimizu, Ishizuka, & Bard, 1997; Simaan & Cruz, 1973). Compared with this, for decision making problems in such as decentralized large firms with divisional independence, it is quite natural to suppose that there exists communication and some cooperative relationship among the decision makers (Sakawa & Nishizaki, 2009).

Assuming that decisions of decision makers in all levels are sequential and all of the decision makers essentially cooperate with each other, Lai (1996) and Shih, Lai, and Lee (1996) proposed solution concepts for two-level linear programming problems. In their methods, the decision makers identify membership functions of the fuzzy goals for their objective functions, and in particular, the decision maker at the upper level also specifies those of the fuzzy goals for the decision variables. The decision maker at the lower level solves a fuzzy programming problem with a constraint with respect to a satisfactory degree of the decision maker at the upper level. Unfortunately, there is a possibility that their method leads a final solution to an undesirable one because of inconsistency between the fuzzy goals of the objective function and those of the decision variables. In order to overcome the problem in their methods, by eliminating the fuzzy goals for the decision variables, Sakawa, Nishizaki, and Uemura (1998, 2000) have proposed interactive fuzzy programming for two-level or multi-level linear programming problems to obtain a satisfactory solution for decision makers. Extensions to two-level linear fractional programming problems (Sakawa, Nishizaki, & Uemura, 2001), decentralized two-level linear programming problems (Sakawa & Nishizaki, 2002; Sakawa, Nishizaki, & Uemura, 2002), two-level linear fractional programming problems with fuzzy parameters (Sakawa et al., 2000), and two-level nonconvex programming problems with fuzzy parameters (Sakawa & Nishizaki, 2002) were provided. Further extensions to two-level linear programming problems with random variables, called stochastic two-level linear programming problems (Sakawa & Katagiri, 2010) and two-level integer

* Corresponding author.

E-mail address: sakawa@hiroshima-u.ac.jp (M. Sakawa).

programming problems (Sakawa, Katagiri, & Matsui, 2012) have also been considered. A recent survey paper of Sakawa and Nishizaki (2012) is devoted to reviewing and classifying the numerous major papers in the area of so-called cooperative multi-level programming.

However, to utilize two-level programming for resolution of conflict in decision making problems in real-world decentralized organizations, it is important to realize that simultaneous considerations of both fuzziness (Sakawa, 1993, 2000, 2001) and randomness (Birge & Louveaux, 1997; Sakawa & Kato, 2008; Stancu-Minasian, 1984) would be required. Fuzzy random variables, first introduced by Kwakernaak (1978), have been developing (Kruse & Meyer, 1987; Liu & Liu, 2003; Puri & Ralescu, 1986), and an overview of the developments of fuzzy random variables was found in Gil, Lopez-Díaz, and Ralescu (2006). Studies on linear programming problems with fuzzy random variable coefficients, called fuzzy random linear programming problems, were initiated by Wang and Qiao (1993) and Qiao, Zhang, and Wang (1994) as seeking the probability distribution of the optimal solution and optimal value. Optimization models for fuzzy random linear programming problems were first developed by Luhandjula (1996) and Luhandjula and Gupta (1996) and further developed by Liu (2001a, 2001b) and Rommelfanger (2007). A brief survey of major fuzzy stochastic programming models including fuzzy random programming was found in the paper by Luhandjula (2006).

Under these circumstances, in this paper, assuming cooperative behavior of the decision makers, we consider solution methods for decision making problems in hierarchical organizations under fuzzy random environments. To deal with the formulated two-level linear programming problems involving fuzzy random variables, α -level sets of fuzzy random variables are introduced and an α -stochastic two-level linear programming problem is defined for guaranteeing the degree of realization of the problem. Taking into account vagueness of judgments of decision makers, fuzzy goals are introduced and the α -stochastic two-level linear programming problem is transformed into the problem to maximize the satisfaction degree for each fuzzy goal. Following probability maximization, the transformed stochastic two-level programming problem can be reduced to a deterministic one. Interactive fuzzy programming to obtain a satisfactory solution for the decision maker at the upper level in consideration of the cooperative relation between decision makers is presented. It is shown that all of the problems to be solved in the proposed interactive fuzzy programming can be easily solved by the simplex method or the combined use of the bisection method and the simplex method.

2. Fuzzy random two-level linear programming problem

Fuzzy random variables, first introduced by Kwakernaak (1978), have been defined in various ways (Kwakernaak, 1978; Kruse & Meyer, 1987; Liu & Liu, 2003; Puri & Ralescu, 1986). For example, as a special case of fuzzy random variables given by Kruse and Meyer (1987) defined a fuzzy random variable as follows.

Definition 1. Fuzzy random variable Let (Ω, B, P) be a probability space, $F(\mathcal{R})$ the set of fuzzy numbers with compact supports and X a measurable mapping $\Omega \rightarrow F(\mathcal{R})$. Then X is a fuzzy random variable if and only if given $\omega \in \Omega$, $X_\alpha(\omega)$ is a random interval for any $\alpha \in (0, 1]$, where $X_\alpha(\omega)$ is an α -level set of the fuzzy set $X(\omega)$.

Although there exist some minor differences in several definitions of fuzzy random variables, fuzzy random variables are considered to be random variables whose observed values are fuzzy sets.

In this paper, we deal with two-level linear programming problems involving fuzzy random variable coefficients in objective functions formulated as:

$$\left. \begin{array}{l} \text{minimize}_{\text{for DM1}} \quad z_1(\mathbf{x}_1, \mathbf{x}_2) = \tilde{\tilde{C}}_{11}\mathbf{x}_1 + \tilde{\tilde{C}}_{12}\mathbf{x}_2 \\ \text{minimize}_{\text{for DM2}} \quad z_2(\mathbf{x}_1, \mathbf{x}_2) = \tilde{\tilde{C}}_{21}\mathbf{x}_1 + \tilde{\tilde{C}}_{22}\mathbf{x}_2 \\ \text{subject to} \quad A_1\mathbf{x}_1 + A_2\mathbf{x}_2 \leq \mathbf{b} \\ \quad \quad \quad \mathbf{x}_1 \geq \mathbf{0}, \mathbf{x}_2 \geq \mathbf{0} \end{array} \right\} \quad (1)$$

where \mathbf{x}_1 is an n_1 dimensional decision variable column vector for the DM at the upper level (DM1), \mathbf{x}_2 is an n_2 dimensional decision variable column vector for the DM at the lower level (DM2), $z_1(\mathbf{x}_1, \mathbf{x}_2)$ is the objective function for DM1 and $z_2(\mathbf{x}_1, \mathbf{x}_2)$ is the objective function for DM2.

It should be emphasized here that randomness and fuzziness of the coefficients are denoted by the “dash above” and “wave above” i.e., “-” and “~”, respectively. In this formulation, \mathbf{x}_1 is an n_1 dimensional decision variable column vector for the DM at the upper level (DM1), \mathbf{x}_2 is an n_2 dimensional decision variable column vector for the DM at the lower level (DM2), $z_1(\mathbf{x}_1, \mathbf{x}_2)$ is the objective function for DM1 and $z_2(\mathbf{x}_1, \mathbf{x}_2)$ is the objective function for DM2. In (1), $\tilde{\tilde{C}}_{lj}$, $l = 1, 2, j = 1, 2$ are vectors whose elements $\tilde{\tilde{C}}_{ljk}$, $k = 1, 2, \dots, n_j$ are fuzzy random variables characterized by the following membership function:

$$\mu_{\tilde{\tilde{C}}_{ljk}}(\tau) = \begin{cases} L\left(\frac{\tilde{d}_{ljk}-\tau}{\beta_{ljk}}\right), & \text{if } \tau \leq \tilde{d}_{ljk} \\ R\left(\frac{\tau-\tilde{d}_{ljk}}{\gamma_{ljk}}\right), & \text{otherwise} \end{cases}$$

where $L: [0, \infty) \rightarrow [0, 1]$ is a monotone decreasing function defined as $L(t) = \max\{0, \lambda(t)\}$ and $R: [0, \infty) \rightarrow [0, 1]$ is also a monotone decreasing function defined as $R(t) = \max\{0, \rho(t)\}$. Here, $\lambda(t)$ and $\rho(t)$ are monotone decreasing functions which satisfy $\lambda(0) = 1$ and $\rho(0) = 1$, respectively. Furthermore, parameters \tilde{d}_{ljk} , β_{ljk} and γ_{ljk} represent a mean value, the left spread and the right spread of $\tilde{\tilde{C}}_{ljk}$, respectively. In this paper, parameters \tilde{d}_{ljk} are random variables defined as $\tilde{d}_{ljk} = d_{ljk}^1 + \tilde{t}_l d_{ljk}^2$, using random variables \tilde{t}_l with mean M_l , $l = 1, 2$. This definition of random variables is one of the simplest randomization modeling of coefficients using dilation and translation of random variables, as discussed by Stancu-Minasian (1990).

Fig. 1 illustrates an example of the membership function of a fuzzy random variable $\tilde{\tilde{C}}_{ljk}$.

Fuzzy random two-level linear programming problems formulated as (1) are often seen in actual decision making situations. For example, consider a supply chain planning (Roghianian, Sadjadi, & Aryanezhad, 2007) where the distribution center (DM1) and the production part (DM2) hope to minimize the distribution cost and




Fig. 1. An example of the membership function $\mu_{\tilde{\tilde{C}}_{ljk}}(\cdot)$ of a fuzzy random variable $\tilde{\tilde{C}}_{ljk}$.

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات