



Mehar's method for solving fully fuzzy linear programming problems with L - R fuzzy parameters



Jagdeep Kaur*, Amit Kumar

School of Mathematics and Computer Applications Thapar University, Patiala 147 004, India

ARTICLE INFO

Article history:

Received 22 May 2012

Received in revised form 16 October 2012

Accepted 24 January 2013

Available online 27 February 2013

Keywords:

Fuzzy linear programming

Fuzzy optimal solution

Ranking function

Fuzzy numbers

ABSTRACT

To the best of our knowledge, there is no method in literature for solving such fully fuzzy linear programming (FLP) problems in which some or all the parameters are represented by unrestricted L - R flat fuzzy numbers. Also, to propose such a method, there is need to find the product of unrestricted L - R flat fuzzy numbers. However, there is no method in the literature to find the product of unrestricted L - R flat fuzzy numbers.

In this paper, firstly the product of unrestricted L - R flat fuzzy numbers is proposed and then with the help of proposed product, a new method (named as Mehar's method) is proposed for solving fully FLP problems. It is also shown that the fully FLP problems which can be solved by the existing methods can also be solved by the Mehar's method. However, such fully FLP problems in which some or all the parameters are represented by unrestricted L - R flat fuzzy numbers can be solved by Mehar's method but can not be solved by any of the existing methods.

© 2013 Elsevier Inc. All rights reserved.

1. Introduction

Linear programming is one of the most frequently applied operation research techniques. Although, it has been investigated and expanded for more than six decades by many researchers and from the various point of views, it is still useful to develop new approaches in order to better fit the real world problems within the framework of linear programming.

In conventional approach, parameters of linear programming models must be well defined and precise. However, in real world environment, this is not a realistic assumption. Usually, the value of many parameters of a linear programming model is estimated by experts. Clearly, it can not be assumed the knowledge of experts is so precise. Since, Bellman and Zadeh [1] proposed the concept of decision making in fuzzy environments, a number of researchers have exhibited their interest to solve the FLP problems [2–8] and fully FLP problems [9–15].

On the basis of deep study of the existing methods for solving fully FLP problems, it can be concluded that there is no method in the literature for solving fully FLP problems in which some or all the parameters are represented by unrestricted L - R flat fuzzy numbers.

This paper is organised as follows: In Section 2, some basic definitions and arithmetic operations of L - R flat fuzzy numbers are presented. In Section 3, limitations of the existing method [9] are pointed out. In Section 4, product of unrestricted L - R flat fuzzy numbers is introduced. In Section 5, a new method, named as Mehar's method, is proposed to find the fuzzy optimal solution of fully FLP problems. In Section 6, advantages of the Mehar's method over the existing methods are discussed and to illustrate the Mehar's method numerical example is solved. Obtained results are discussed in Section 7. Conclusions are discussed in Section 8.

* Corresponding author.

E-mail addresses: deepi.thaparian@gmail.com (J. Kaur), amit_rs_iitr@yahoo.com (A. Kumar).

2. Preliminaries

In this section, some basic definitions and arithmetic operations of *L-R* flat fuzzy numbers are presented.

2.1. Basic definitions

In this section, some basic definitions of *L-R* flat fuzzy numbers are presented.

Definition 2.1 [16]. A function $L : [0, \infty) \rightarrow [0, 1]$ (or $R : [0, \infty) \rightarrow [0, 1]$) is said to be reference function of fuzzy number if and only if

- (i) $L(0) = 1$ (or $R(0) = 1$)
- (ii) L (or R) is non-increasing on $[0, \infty)$.

Definition 2.2 [16]. A fuzzy number \tilde{A} , defined on universal set of real numbers \mathbb{R} , denoted as $(m, n, \alpha, \beta)_{LR}$, is said to be an *LR* flat fuzzy number if its membership function $\mu_{\tilde{A}}(x)$ is given by

$$\mu_{\tilde{A}}(x) = \begin{cases} L\left(\frac{m-x}{\alpha}\right) & x \leq m, \alpha > 0, \\ R\left(\frac{x-n}{\beta}\right) & x \geq n, \beta > 0, \\ 1 & m \leq x \leq n. \end{cases}$$

Definition 2.3 [14]. An *L-R* flat fuzzy number $\tilde{A} = (m, n, \alpha, \beta)_{LR}$ is said to be non-negative *L-R* flat fuzzy number if $m - \alpha \geq 0$ and is said to be non-positive *L-R* flat fuzzy number if $n + \beta \leq 0$.

Definition 2.4 [14]. An *L-R* flat fuzzy number $\tilde{A} = (m, n, \alpha, \beta)_{LR}$ is said to be unrestricted *L-R* flat fuzzy number if $m - \alpha$ is a real number.

Definition 2.5 [16]. Let $\tilde{A} = (m, n, \alpha, \beta)_{LR}$ be an *L-R* flat fuzzy number and λ be a real number in the interval $[0, 1]$ then the crisp set, $A_\lambda = \{x \in X : \mu_{\tilde{A}}(x) \geq \lambda\} = [m - \alpha L^{-1}(\lambda), n + \beta R^{-1}(\lambda)]$, is said to be λ -cut of \tilde{A} .

Definition 2.6 [16]. Let $\tilde{A}_1 = (m_1, n_1, \alpha_1, \beta_1)_{LR}$ and $\tilde{A}_2 = (m_2, n_2, \alpha_2, \beta_2)_{LR}$, be any *L-R* flat fuzzy numbers then $\tilde{A}_1 = \tilde{A}_2$ if $m_1 = m_2, n_1 = n_2, \alpha_1 = \alpha_2$ and $\beta_1 = \beta_2$.

2.2. Arithmetic operations

In this section, the arithmetic operations between *L-R* flat fuzzy numbers are presented [16].

Let $\tilde{A}_1 = (m_1, n_1, \alpha_1, \beta_1)_{LR}$, $\tilde{A}_2 = (m_2, n_2, \alpha_2, \beta_2)_{LR}$ be any *L-R* flat fuzzy numbers and $\tilde{A}_3 = (m_3, n_3, \alpha_3, \beta_3)_{RL}$ be any *R-L* flat fuzzy number. Then,

- (i) $\tilde{A}_1 \oplus \tilde{A}_2 = (m_1 + m_2, n_1 + n_2, \alpha_1 + \alpha_2, \beta_1 + \beta_2)_{LR}$
- (ii) $\tilde{A}_1 \ominus \tilde{A}_3 = (m_1 - n_3, n_1 - m_3, \alpha_1 + \beta_3, \beta_1 + \alpha_3)_{LR}$
- (iii) If \tilde{A}_1 and \tilde{A}_2 both are non-negative, then $\tilde{A}_1 \odot \tilde{A}_2 \simeq (m_1 m_2, n_1 n_2, m_1 \alpha_2 + \alpha_1 m_2 - \alpha_1 \alpha_2, n_1 \beta_2 + \beta_1 n_2 + \beta_1 \beta_2)_{LR}$
- (iv) If \tilde{A}_1 is non-positive and \tilde{A}_2 is non-negative, then $\tilde{A}_1 \odot \tilde{A}_2 \simeq (m_1 n_2, n_1 m_2, \alpha_1 n_2 - m_1 \beta_2 + \alpha_1 \beta_2, \beta_1 m_2 - n_1 \alpha_2 - \beta_1 \alpha_2)_{LR}$
- (v) If \tilde{A}_1 is non-negative and \tilde{A}_2 is non-positive, then $\tilde{A}_1 \odot \tilde{A}_2 \simeq (n_1 m_2, m_1 n_2, n_1 \alpha_2 - \beta_1 m_2 + \beta_1 \alpha_2, m_1 \beta_2 - \alpha_1 n_2 - \alpha_1 \beta_2)_{LR}$
- (vi) If \tilde{A}_1 and \tilde{A}_2 both are non-positive, then $\tilde{A}_1 \odot \tilde{A}_2 \simeq (n_1 n_2, m_1 m_2, -n_1 \beta_2 - \beta_1 n_2 - \beta_1 \beta_2, -m_1 \alpha_2 - \alpha_1 m_2 + \alpha_1 \alpha_2)_{LR}$
- (vii) $\lambda \tilde{A}_1 = \begin{cases} (\lambda m_1, \lambda n_1, \lambda \alpha_1, \lambda \beta_1)_{LR} & \lambda \geq 0 \\ (\lambda n_1, \lambda m_1, -\lambda \beta_1, -\lambda \alpha_1)_{RL} & \lambda < 0 \end{cases}$

There also exist another formula [16] for the product of such *L-R* flat fuzzy numbers in which the spreads are smaller as compared to the mean values:

- (i) If \tilde{A}_1 and \tilde{A}_2 both are non-negative, then $\tilde{A}_1 \otimes \tilde{A}_2 \simeq (m_1 m_2, n_1 n_2, m_1 \alpha_2 + \alpha_1 m_2, n_1 \beta_2 + \beta_1 n_2)_{LR}$

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات