



Sensitivity Analysis in Periodic Matrix Models: A Postscript to Caswell and Trevisan[†]

M. LESNOFF*

Centre de Coopération Internationale en Recherche Agronomique pour le
Développement (CIRAD – EMVT) TA/30A
34398 Montpellier Cedex 5, France
and

International Livestock Research Institute (ILRI)
P.O. Box 5689, Addis Ababa, Ethiopia
m.lesnoff@cgiar.org

P. EZANNO

Centre de Coopération Internationale en Recherche Agronomique pour le
Développement (CIRAD – EMVT) TA/30A
34398 Montpellier Cedex 5, France
pauline.ezanno@cirad.fr

H. CASWELL

Biology Department, MS #34
Woods Hole Oceanographic Institution
Woods Hole, MA 02543-1049, U.S.A.
hcaswell@whoi.edu

(Received April 2002; revised and accepted December 2002)

Abstract—Periodic matrix population models are a useful approach to modelling cyclic variations in demographic rates. Caswell and Trevisan [1] introduced the perturbation analysis (sensitivities and elasticities) of the per-cycle population growth rate for such models. Although powerful, their method can be time-consuming when the dimension of the matrices is large or when cycles are composed of many phases. We present a more efficient method, based on a very simple matrix product. We compared the two methods for matrices of different sizes. We observed a reduction in calculation time on the order of 24% with the new method for a set of 26 within-year Leslie matrices of size 287×287 . The time saving may become particularly significant when sensitivities are used in Monte Carlo or bootstrap simulations. © 2003 Elsevier Science Ltd. All rights reserved.

Keywords—Sensitivity, Elasticity, Periodic matrix models, Population dynamics, Population growth rate.

Periodic matrix population models are a useful approach to modelling cyclic variations in demographic rates, such as are caused by seasonality within the year or by interannual cyclic variability. See [2, Chapter 13] for a review of biological applications. Caswell and Trevisan [1]

*Author to whom all correspondence should be addressed. Please send to author's second address.

†See [1].

introduced the perturbation analysis (sensitivities and elasticities) of population growth rate for periodic models; the objective of our postscript is to introduce a simpler way to calculate these sensitivities and elasticities.

We suppose here that the cycle is composed of K “phases” (e.g., a year composed of $K = 4$ seasons, or of $K = 26$ two-week phases). The phases need not be of the same duration. The matrices $\mathbf{B}_1, \mathbf{B}_2, \dots, \mathbf{B}_K$ denote the population projection matrices for the different phases. That is, matrix \mathbf{B}_i projects the population from phase i to phase $i + 1$; the phases are cyclic, so that \mathbf{B}_K projects the population from phase K back to phase 1. The starting point of the cycle is arbitrary. Consider a cycle starting at the beginning of phase k and let $\mathbf{x}(t)$ denote the population state vector at time t . The dynamics over the whole cycle are given by [1]

$$\mathbf{x}(t + K) = \mathbf{B}_{k-1}\mathbf{B}_{k-2} \dots \mathbf{B}_1\mathbf{B}_K\mathbf{B}_{K-1} \dots \mathbf{B}_k\mathbf{x}(t), \tag{1}$$

$$\equiv \mathbf{A}_k\mathbf{x}(t). \tag{2}$$

The asymptotic properties of such models have been described in Skellam [3] and Caswell [1,2,4]. Under weak conditions of primitivity, the asymptotic population growth rate λ (on the per-cycle scale) is the common dominant eigenvalue of the product-matrices \mathbf{A}_k (all the \mathbf{A}_k have the same eigenvalues).

Our concern here is to calculate the sensitivities of λ to changes in the entries of each of the matrices \mathbf{B}_k . Using the notation in [1], let $a_{ij}^{(k)}$ denote the (i, j) entry of the product-matrix \mathbf{A}_k and S_{A_k} the sensitivity matrix of \mathbf{A}_k , i.e., the matrix whose (i, j) entry is the partial derivative $\frac{\partial \lambda}{\partial a_{ij}^{(k)}}$. This matrix can be calculated directly from the eigenvectors of \mathbf{A}_k , but because the entries of \mathbf{A}_k are complicated combinations of the phase-specific demographic rates, these sensitivities are difficult to interpret. Of more interest is the sensitivity matrix S_{B_k} (the matrix whose (i, j) entry is the partial derivative $\frac{\partial \lambda}{\partial b_{ij}^{(k)}}$ where $b_{ij}^{(k)}$ denotes the (i, j) entry of \mathbf{B}_k). Caswell and Trevisan [1] showed that these sensitivity matrices are given by

$$\mathbf{S}_{B_k} = (\mathbf{B}_{k-1}\mathbf{B}_{k-2} \dots \mathbf{B}_1\mathbf{B}_K\mathbf{B}_{K-1} \dots \mathbf{B}_{k+1})^T \mathbf{S}_{A_k} \quad k = 1, \dots, K. \tag{3}$$

Equation (3) is powerful and easy to implement in appropriate software. However, it requires the calculation of K sensitivity matrices \mathbf{S}_{A_k} . This calculation could become time-consuming when the dimension of matrices \mathbf{B}_k is large and when there are many phases in the annual cycle. Next, we present a more efficient method.

Since the sensitivity $\frac{\partial \lambda}{\partial b_{ij}^{(k)}}$ is independent of which cyclic permutation of the \mathbf{B} matrices is considered, we suppose here for notational simplicity, and without loss of generality, that the cyclic projection matrix is $\mathbf{A}_1 = \mathbf{B}_K\mathbf{B}_{K-1} \dots \mathbf{B}_1 \equiv \mathbf{A}$. The population growth rate λ can be seen as a composite function of the variables a_{mn} and $b_{ij}^{(k)}$, i.e.,

$$\lambda = \lambda \left(a_{mn} \left(b_{ij}^{(k)} \right) \right), \quad i, j, m, n = 1, \dots, q; \quad k = 1, \dots, K, \tag{4}$$

where q is the dimension of matrices \mathbf{A}_k and \mathbf{B}_k .

From the chain rule, the partial derivative of λ with respect to $b_{ij}^{(k)}$ is

$$\frac{\partial \lambda}{\partial b_{ij}^{(k)}} = \sum_{m,n} \frac{\partial \lambda}{\partial a_{mn}} \frac{\partial a_{mn}}{\partial b_{ij}^{(k)}}. \tag{5}$$

Our problem is to find the derivatives $\frac{\partial a_{mn}}{\partial b_{ij}^{(k)}}$ in a more efficient way than that of Caswell and Trevisan [1]. To do so, we rewrite matrix \mathbf{A} as

$$\mathbf{A} = \mathbf{C}\mathbf{B}_k\mathbf{G}, \tag{6}$$

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات