




ELSEVIER

Contents lists available at ScienceDirect

Journal of Obsessive-Compulsive and Related Disorders

journal homepage: www.elsevier.com/locate/jocrd

Part 3. A question of perspective: The association between intrusive thoughts and obsessionality in 11 countries



David A. Clark^{a,*}, Jon Abramowitz^b, Gillian M. Alcolado^c, Pino Alonso^d, Amparo Belloch^e,
Martine Bouvard^f, Meredith E. Coles^g, Guy Doron^h, Hector Fernández-Álvarezⁱ,
Gemma Garcia-Soriano^e, Marta Ghisi^j, Beatriz Gomezⁱ, Mujgan Inozu^k,
Richard Moulding^{l,m}, Adam S. Radomsky^c, Giti Shamsⁿ, Claudio Sica^o,
Gregoris Simos^p, Wing Wong^q

^a Department of Psychology, University of New Brunswick, Fredericton, NB, Canada E3B 9P8^b Department of Psychology, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA^c Department of Psychology, Concordia University, Montreal, QC, Canada^d Department of Psychiatry, Bellvitge Hospital, Barcelona, Spain^e Department of Personality Psychology, University of Valencia, Valencia, Spain^f Department of Psychology, University of Savoie, Chambéry, France^g Department of Psychology, Binghamton University, Binghamton, NY, USA^h Department of Psychology, Interdisciplinary Center, Herzliya, Israelⁱ Aigle Foundation, Buenos Aires, Argentina^j Department of General Psychology, University of Padova, Padova, Italy^k Department of Psychology, Abant İzzet Baysal University, Bolu, Turkey^l Department of Psychology, Deakin University, Melbourne, VIC, Australia^m Brain and Psychological Sciences Research Centre, Swinburne University of Technology, Melbourne, VIC, Australiaⁿ Department of Psychiatry, Tehran University, Medical Science, Tehran, Iran^o Department of Human Health Science, University of Firenze, Firenze, Italy^p Department of Educational and Social Policy, University of Macedonia, Thessaloniki, Greece^q Departments of Psychiatry and Psychology, Chinese University of Hong Kong, Shatin, Hong Kong

ARTICLE INFO

Article history:

Received 24 June 2013

Received in revised form

6 December 2013

Accepted 9 December 2013

Available online 17 December 2013

Keywords:

OCD

Obsessions

Intrusive thoughts

Obsessionality

Obsessive-compulsive inventory-revised

ABSTRACT

A key assumption of contemporary cognitive-behavioral models of obsessive-compulsive disorder (OCD) is that obsessional thoughts exist on a continuum with “normal” unwanted intrusive thoughts. Recently, however, some authors have challenged this notion. The present study aimed to clarify (a) the extent that different types of intrusive thoughts in nonclinical individuals are associated with obsessionality, (b) the relative contribution of frequency, distress and control ratings to obsessionality, and (c) the extent that existing findings (primarily from North American or European samples) generalize to other countries in the world. Five hundred and fifty-four non clinical individuals from 11 different countries were administered an interview assessing the presence, frequency, distress, and perceived control of different types of intrusive thoughts. Participants also completed measures of obsessional beliefs, obsessive-compulsive (OC) symptoms, and depression. Results from data analyses supported the universality of unwanted intrusive thoughts, the continuity of normal and abnormal obsessions, and the specificity of dirt/contamination, doubt and miscellaneous intrusions to OC symptoms. Implications for intrusive thoughts as a potential vulnerability factor for OCD are discussed.

© 2013 Elsevier Ltd. All rights reserved.

1. Introduction

Cognitive-behavioral researchers and practitioners widely believe that obsessions arise from the normal unwanted intrusive thoughts,

images or impulses that are universally experienced by all individuals. Born out of the seminal study by [Rachman and de Silva \(1978\)](#), showing that the majority of individuals experience occasional unwanted intrusive thoughts that are indistinguishable from the content of clinical obsessions, subsequent cognitive-behavioral treatment (CBT) models of obsessive-compulsive disorder (OCD) have embraced the assumption of continuity between “normal” and abnormal obsessions ([Clark, 2004](#); [Freeston, Rhéaume, & Ladouceur, 1996](#); [Rachman, 1997, 1998, 2003](#); [Rachman & Hodgson, 1980](#);

* Correspondence to: Department of Psychology, University of New Brunswick, PO Box 4400, Fredericton, New Brunswick Canada E3B 5A3.
E-mail address: clark@unb.ca (D.A. Clark).

Salkovskis, 1985, 1989a). CBT models contend that any normal intrusive thought can escalate into an obsession if it is erroneously interpreted as signifying a serious personal threat to self or significant others and the person attempts to reduce or neutralize the distressing qualities of the intrusion by engaging in some futile behavioral or mental activity (Rachman, 1997, 1998). From this perspective, any thought, regardless of content or context, could morph into a clinical obsession if associated with pathological appraisal and neutralization processes (Salkovskis & Freeston, 2001).

Most of the published research on intrusive thoughts has been conducted on western samples and there have been no large scale investigations of intrusive thoughts across a diverse range of nationalities. Thus the universality of unwanted intrusions, their relationship to OC symptoms and, by extension, the cross-cultural relevance of the CBT model of OCD has not been tested in a multinational comparative study. Like the other papers in this special issue, a major objective of the current study is to determine the cross-cultural generalizability of the intrusion/OC symptom relationship in a large multinational comparative study.

Not all OCD researchers readily embrace the continuity of normal and abnormal obsessions, or the universality of obsessive-like intrusive thoughts. Prior to the Rachman and de Silva (1978) research, the prevailing notion in psychiatry was that obsessions represented aberrant thinking that was distinct from normal cognition (Black, 1974; Lewis, 1936; Pollitt, 1957). The view that clinical obsessions are qualitatively distinct from normal cognition is more consistent with the categorical approach to psychiatric nosology represented in DSM-5 than is the continuity assumption of obsessions (American Psychiatric Association, 2013). Moreover, several recent empirical studies have challenged the continuity assumption. In their review of the intrusive thoughts literature, Julien, O'Connor, and Aardema (2007) concluded that intrusive thoughts do not always exhibit a stronger correlation with OCD symptoms than with other anxiety symptoms or depression, nor do the phenomena show the expected distinctiveness from other types of negative cognitions such as worry and negative automatic thoughts. In support of this argument, Rassin and Muris (2006), Rassin, Cogle, and Muris (2007) reported on two studies that appear to indicate that some "abnormal" intrusions have a closer relationship to obsessionality than do other more normal intrusions, with non-clinical participants reporting more of the latter than the former. This finding challenges the assumption of continuity between nonclinical and clinical intrusions.

There are, however, two difficulties with the conclusions reached in these studies. First, there was a very strong correlation between frequency of "normal" and "abnormal" intrusive thoughts ($r = .80$), indicating that people who tended to endorse one type of thought also endorsed the other. Second, their list of "normal" and "abnormal" intrusions may have low content validity because the list of "normal intrusions" included many instances of angry, worrisome, and generally negative automatic thoughts (e.g., "they shouldn't do that"). Thus it is not surprising that the Rassin et al. findings are inconsistent with those of Lee, Lee, Kim, Kwon, and Telch (2005), who found that the more realistic reactive intrusions, which are similar to Rassin and Muris' (2006) "normal obsession" content, had a stronger relation to obsessive compulsive (OC) symptoms than the bizarre, highly aversive and unrealistic "auto-genous" intrusions, which are like Rassin and Muris' abnormal obsessions category.

There have now been several replication studies documenting the almost unanimous presence of unwanted intrusive thoughts in healthy, nonclinical samples (Freeston, Ladouceur, Thibodeau, & Gagnon, 1991; Parkinson & Rachman, 1981; Salkovskis & Harrison, 1984), even when more stringent criteria of intrusions are utilized (Belloch, Morillo, Lucero, Cabedo, & Carrió, 2004; Lee & Kwon, 2003; Purdon & Clark, 1993). Radomsky et al. (2014) found that across 13 countries the vast majority of nonclinical individuals

(93.6%) reported experiencing unwanted intrusive thoughts with obsessional content within the past three months, although some differences were evident between countries. These findings, then, support one of the basic assertions of the cognitive-behavioral model of OCD; that unwanted obsessive-like intrusive thoughts are highly prevalent in the nonclinical population, and this phenomenon may be present regardless of nationality, religion or culture.

A second major premise of the cognitive-behavioral perspective is that the frequency, appraisal and control of intrusive thoughts will exhibit a specific relationship with obsessional symptoms. The most stringent version of this hypothesis argues for a causal relation between intrusions and obsessional symptoms. Although most studies, including the present, are cross-sectional in nature and so cannot investigate causal relations, these studies have been able to test a precondition of causality; that is, whether unwanted intrusive thoughts have a unique and significant relationship with obsessionality. Various studies report that the frequency of intrusive thoughts has a stronger association with obsessional than with anxious or depressive symptoms (Barrera & Norton, 2011; García-Soriano, Belloch, Morillo, & Clark, 2011; Lee et al., 2005; Purdon & Clark, 1993), although failure to find specificity has also been reported (Clark, 1992). Other studies find that it is not the frequency of intrusive thoughts that is uniquely associated with OC symptoms, but rather how the thought is appraised or evaluated (Corcoran & Woody, 2008; Freeston & Ladouceur, 1993; Freeston, Ladouceur, Thibodeau, & Gagnon, 1992). Still other studies report that it is perceived uncontrollability or effort to control intrusions that is specific to obsessionality (Grisham & Williams, 2009; Morillo, Belloch, & Garcia-Soriano, 2007; see Magee, Harden, & Teachman, 2012 for further discussion).

To date several fundamental issues about the relation between unwanted intrusive thoughts and obsessional symptoms remain unresolved. First, it is not clear whether the content of intrusive thoughts is entirely irrelevant to OC symptoms. It may be that some types of intrusions have a closer association with or relevance to obsessionality than do others, but answers to this question have been hampered by studies that use small, relatively homogenous samples and rely on retrospective self-report intrusion and symptom measures that have high levels of shared method variance. Second although the most consistent finding is that frequency of unwanted intrusions is related to OC symptoms, the relative contribution of appraisal and control efforts is still not well understood. Finally, much of the work on intrusions and obsessionality is based on North American or European samples. It is unknown whether these relationships generalize to populations in other regions of the world. Nevertheless, based on the research to date, we can expect that unwanted intrusive thoughts with varying content will have a specific relationship to OC symptoms across all cultures and that perceived distress and control of intrusions will have a significant relationship with OC symptoms even after controlling for the frequency of intrusive thoughts.

To determine content-specificity, generalizability across cultures and contribution of appraisals to the intrusion/OC symptom relationship, the present study focuses on analyses conducted on the frequency, distress, importance of control, and difficulty of control ratings completed for the seven types of intrusive thoughts assessed in the International Intrusive Thoughts Interview Schedule (IITIS; RCIF, 2007). Between-group comparisons across country as well as a series of hierarchical regression analyses investigated the specific relation between frequency, distress and perceived control of unwanted intrusions and distress ratings for OCD symptoms. It was hypothesized that frequency of all seven types of intrusions would have a significantly greater association with OC than with depressive symptoms, and that frequency of intrusions would be

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات