



PERGAMON

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Journal of Behavior Therapy
and Experimental Psychiatry 34 (2003) 161–170

JOURNAL OF
behavior
therapy
and
experimental
psychiatry

www.elsevier.com/locate/jbtep

Emotional responding to hyperventilation as a predictor of agoraphobia status among individuals suffering from panic disorder

Michael J. Telch^{a,*}, Kristine Jacquin^b, Jasper A.J. Smits^a,
Mark B. Powers^a

^a *Laboratory for the Study of Anxiety Disorders, Department of Psychology, University of Texas at Austin, Mezes 330, Mail Stop B3800, Austin, TX 78712, USA*

^b *Department of Psychology, Mississippi State University, USA*

Received 5 July 2002; received in revised form 6 December 2002; accepted 6 May 2003

Abstract

Some data suggest that panic patients with extensive agoraphobia (PDA) display more intense respiratory distress during their panic attacks than Panic disorder (PD) patients. However, no studies have determined if PDA patients also show heightened sensitivity to a respiratory challenge compared to PD patients. The current study examined the differential emotional responding to hyperventilation among PDA patients, PD patients, and a non-clinical group with a history of panic attacks. Response to hyperventilation challenge did not distinguish non-clinical panickers from panic patients; however, behavioral tolerance to hyperventilation challenge significantly predicted agoraphobia status among panic disorder patients, even after controlling for demographic and clinical status variables.

© 2003 Elsevier Ltd. All rights reserved.

Keywords: Panic disorder; Agoraphobia; Hyperventilation; Respiration; Anxiety sensitivity; Challenge test

1. Introduction

Challenging patients in the laboratory has been a widely used tool in the study of panic disorder (PD). There is now ample evidence demonstrating that compared to normal controls, PD patients display heightened physiologic activation, marked rise

*Corresponding author. Tel.: +1-512-471-3393; fax: +1-512-471-6175.

E-mail address: telch@forum.utexas.edu (M.J. Telch).

in fear, and increased probability of panic in response to voluntary hyperventilation (Beck & Scott, 1988; Gorman et al., 1984, 1988, 1994; Holt & Andrews, 1989a, b; Nardi, Valenca, Nascimento, Mezzasalma, & Zin, 1999, 2001; Papp et al., 1997), inhalation of 5% CO₂ (Gorman et al., 1984, 1988, 1994; Papp et al., 1997; Woods, Charney, Goodman, & Heninger, 1988), 7% CO₂ (Gorman et al., 1994; Papp et al., 1997), and 35% CO₂ (Gorman et al., 1990; Griez, Lousberg, van den Hout, & van der Molen, 1987; Papp et al., 1993; Perna et al., 1995, 1994; Perna, Gabriele, Caldirola, & Bellodi, 1995). Moreover, PD sufferers show greater sensitivity to 5% and 35% CO₂ than patients with obsessive compulsive disorder (Griez, de Loof, Pols, Zandbergen, & Lousberg, 1990; Perna, Bertani, Arancio, Ronchi, & Bellodi, 1995), generalized anxiety disorder (Perna, Bussi, Allevi, & Bellodi, 1999; Verburg, Griez, Meijer, & Pols, 1995) or depression (Kent et al., 2001; Perna, Barbini, Cocchi, Bertani, & Gasperini, 1995) and greater sensitivity to hyperventilation challenges compared to those with generalized anxiety disorder, or social phobia (Holt & Andrews, 1989b; Rapee, Brown, Antony, & Barlow, 1992). Based on these findings, it has been suggested that respiratory abnormalities play a central role in the pathogenesis of panic disorder (Klein, 1993).

Despite the plethora of respiratory challenge studies, none to our knowledge have examined differential emotional responding to hyperventilation among panic patients with and without extensive agoraphobic avoidance. When present in panic disorder, agoraphobia is associated with markedly greater disruption in quality of life (de Jong & Bouman, 1995), significantly higher rates of comorbid depression (Goisman et al., 1994), substance abuse (Rapee & Medoro, 1994), and a less favorable response to both pharmacologic and cognitive-behavioral treatment (Goisman et al., 1994; Keller et al., 1994). Although panic patients with and without agoraphobia do not differ on panic attack frequency (Cox, Endler, & Swinson, 1995; Craske & Barlow, 1988; Rapee & Murrell, 1988; Telch, Brouillard, Telch, Agras, & Taylor, 1989), there are some data suggesting that panic patients with agoraphobia may display more intense respiratory distress such as faintness or dizziness during their panic attacks (de Jong & Bouman, 1995; Noyes, Clancy, Garvey, & Anderson, 1987; Telch et al., 1989).

The overall aim of the present investigation was to examine the role of respiratory sensitivity in panic disorder and agoraphobia by comparing emotional responding to voluntary hyperventilation challenge among three panic groups: non-patients with current panic attacks (non-clinical panickers (NP)), patients with PD but minimal or no agoraphobia, and patients with PD plus moderate to severe agoraphobia. Several specific questions were addressed: (a) Do patients with PD (with or without agoraphobia) display greater anxious responding to a respiratory challenge compared to NP? (b) Do patients with PD with significant agoraphobia display heightened emotional responding relative to PD patients with minimal or no agoraphobia? and (c) Do measures of emotional sensitivity to hyperventilation challenge predict patients' clinical status after controlling for clinical and demographic differences. Similarly, do indices of respiratory sensitivity predict agoraphobia status, even after controlling for differences in demographic and clinical characteristics?

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات