

Original articles

Evaluation of the structure of Brazilian State-Trait Anxiety Inventory using a Rasch psychometric approach ^{☆,☆☆}

Márcia Balle Kaipper^{a,1}, Eduardo Chachamovich^{b,1}, Maria Paz Loayza Hidalgo^{a,d},
Iraci Lucena da Silva Torres^{a,c}, Wolnei Caumo^{a,c,e,*}

^aPost-Graduate Program in Medical Sciences, School of Medicine, Universidade Federal do Rio Grande do Sul (UFRGS), Brazil

^bPost-Graduate Program in Psychiatry, School of Medicine, UFRGS, Brazil

^cPharmacology Department, Institute of Basic Health Science of UFRGS, Brazil

^dDepartment of Psychiatry, School of Medicine, UFRGS; Psychiatric Service of Hospital de Clínicas de Porto Alegre (HCPA), Brazil

^ePain and Palliative Care Service, HCPA/UFRGS; Porto Alegre, Brazil

Received 20 August 2008; received in revised form 22 September 2009; accepted 22 September 2009

Abstract

Objective: This study evaluates the State-Trait Anxiety Inventory (STAI) structure using a Rasch psychometric approach, and a refined and shorter STAI version is proposed. **Methods:** A cross-sectional study was performed with 900 inpatients scheduled for elective surgery. Age varied from 18 to 60 years (American Society of Anesthesiologists physical status I–III). Demographic information was collected using a structured questionnaire. The measuring instrument (the STAI) was applied to all patients in the afternoon before the surgery and prior to the patients receiving preoperative sedatives. **Results:** Rasch analysis of the state and trait anxiety scales was performed separately. This analysis demonstrated that the original format of state and trait scales fails to show invariance across the trait-state anxiety level, which results in the unstable performance of items. The refined scale was retested in two subsequent random samples of 300 subjects each, and the results were confirmed. The performance was adequate regardless of gender. In the analysis, some items of the state scale

(items 3,4,9,10,12,15, and 20) were deleted due to poor fit statistics. The remaining 13 items showed unidimensionality, local independence, and adequate index of internal consistency. Also, the original trait scale displayed several weaknesses. First, the four-point Likert response scale proved to be inadequate, and threshold disorders were found in all 20 items. Also, the original trait scale showed insufficient item-trait interaction and several individual item misfits. Following the rescoring process, and retesting in a second random sample, items were excluded (namely Items 3, 4, 11, 13, 14, 15, 18, and 19). The refined version showed local independence, unidimensionality, and adequate fit statistics. **Discussion:** The results indicate that the application of the Rasch model led to the refinement of the classic STAI state and trait scales. In addition, they suggest that these shorter versions have a more suitable psychometric performance and are free of threshold disorders and differential item functioning problems. Published by Elsevier Inc.

Keywords: STAI; Anxiety; Rasch analysis; Surgery

[☆] Conflict of Interest: There was no financial relationship between any of the authors or any commercial interest in the outcome of this study.

^{☆☆} Financial support: This research was supported by the Graduate Research Group (GPPG) at Hospital de Clínicas de Porto Alegre, Brazil and the Postgraduate Program in Medical Sciences, School of Medicine, UFRGS.

* Corresponding author. Coronel Corte Real, 295, Bairro Petropolis, CEP: 90630-008 – Porto Alegre, RS, Brazil. Tel.: +55 51 30237448; fax: +55 51 3339733.

E-mail address: caumo@cpovo.net (W. Caumo).

¹ Both are the first authors.

Introduction

Most patients awaiting elective surgery experience preoperative anxiety [1–4]. The incidence of preoperative anxiety has been reported to range from 11% to 80% among adult patients [5]. This widely range may be explained in part by the assessment methods.

The State-Trait Anxiety Inventory (STAI) is the gold standard for anxiety evaluation used in more than one thousand peer-reviewed studies [6]. The STAI-Form X was published in 1970 [7], and its revision began in 1979. The STAI comprises separate self-report scales for measuring the state (anxiety refers to acute situation-driven episodes and fluctuates over time), and the trait anxiety (a lifelong pattern of anxiety, a personality feature) [8–10]. The S-Anxiety Scale (STAI-S) consists of twenty statements that evaluate how the respondent feels “right now,” at this moment. The T-Anxiety Scale (STAI-T) consists in twenty statements that evaluate how the respondent feels “generally”. The refined version of the STAI-Form X changed around 25% of the questions and it was denominated STAI Form Y (1983) [8]. However, the STAI Form Y persists with the 20 multiple-choice questions for state anxiety and 20 questions for trait anxiety and its structure was based in the classic psychometric parameters. Additionally, it is important to emphasize that even so being the STAI Form Y, the STAI-Form X persists the instrument more widely used to assess state-trait anxiety [5,11]. Thus, it is essential to examine more in depth the STAI-Form X to determine the effectiveness of interventions directed at reducing such anxiety. This is especially true because previous reports have demonstrated that anesthesiologists appear to be inaccurate in assessing patient anxiety during the preoperative visit [1,12]. This moment of the perioperative period is when the patient’s anxiety should be predicted with higher accuracy to allow the identification of patients with high anxiety, so that appropriate steps can be taken to alleviate this anxiety.

Given the importance of studying anxiety across cultures and the need to establish cross-cultural equivalence of instruments, it is important to investigate the psychometric properties of the STAI using a statistical approach, such as Rasch analysis, to investigate the structure of the STAI more in depth. It is important for theory and research, since an instrument could assess anxiety appropriately independent of the situation in which it is experienced or of the interpretation of its meaning. The Rasch measurement theory is a modern psychometric approach for the development and validation of instruments. It has emerged as a powerful tool for examining the performance of an instrument in depth, allowing both the instrument as whole and individual items to be assessed. In addition, the Rasch model is also helpful in providing potential solutions for misperforming instruments. It is suggested that combining both traditional and modern psychometric approaches is a fruitful strategy to enhance the power of validation processes [13]. Furthermore, the use of the Rasch measurement model

for the development and application of mental health instruments has been increasingly stressed [14,15].

The present paper evaluates the STAI-Form X structure using a Rasch psychometric approach, and a refined and shorter STAI-Form X version is proposed. In addition, the discriminate properties of the original STAI-Form X and their reduced forms were used to identify patients more disposed to present higher postoperative pain.

Methods

A cross-sectional study of adult patients was performed in a tertiary care hospital in the city of Porto Alegre, Brazil, after institutional approval by the local ethics committee. Written informed consent was obtained from all patients. The sample was composed of patients admitted to the hospital one day prior to elective surgery with general anesthesia or neural blockage. Patients were classified in terms of physical status according to the American Society of Anesthesiologists criteria (ASA, class status I-III). Subjects were at least 18 years old. Exclusion criteria included a medical history of organic brain damage, mental retardation, difficulty in understanding verbal commands, use of pre-anesthetic medications before the evaluation carried out for the present study, and ophthalmologic or cardiac surgery. All patients were scheduled to be submitted to a variety of elective surgeries classified as minor, medium, and major according to blood loss, degree of pain, invasiveness, degree of monitoring required, and length of stay in the hospital due to the surgical procedure [16,17]. One subgroup of 70 patients who underwent abdominal hysterectomy, were previously included in a randomized clinical trial, and were placebo-treated was used to assess the discriminative properties of the original STAI-Form X and their reduced versions to identify patients more susceptible to report higher levels of postoperative pain. These patients received epidural analgesia with ropivacaine, and after being transferred to the post-anesthesia care unit, they were connected to a morphine patient-controlled analgesia (PCA) pump.

Measurement of anxiety

The STAI-Form X, validated and adapted to Brazilian Portuguese [18–21], was utilized to measure anxiety. The inventory contains two separate lists of 20 items each of self-report rating scales for measuring trait and state anxiety. In responding to the S-STAI-Form X scales, subjects chose the number that best describes the intensity of their feelings: (1) not at all, (2) somewhat, (3) moderately, (4) and very much so. In responding to the T-Anxiety Scale, subjects rate the frequency of their feelings on the following four-point scale: (1) almost never, (2) sometimes, (3) often, and (4) almost always. Each STAI item is given a weighted score of 1 to 4. The total score for each scale ranges from 20 to 80. Higher scores denote higher levels of anxiety.

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات