



ELSEVIER

Journal of Fluency Disorders 30 (2005) 163–188

Journal of
**FLUENCY
DISORDERS**

Evidence-based practice in stuttering: Some questions to consider

Nan Bernstein Ratner*

*Department of Hearing and Speech Sciences, The University of Maryland,
0100 Lefrak Hall, College Park, MD 20742, USA*

Received 30 March 2004; received in revised form 19 April 2005; accepted 22 April 2005

Abstract

A recent forum in *JFD* (28/3, 2003) evaluated the status of evidence-based practice in fluency disorders, and offered recommendations for improvement. This article re-evaluates the level of support available for some popular approaches to stuttering therapy and questions the relative value placed on some types of programs endorsed by the forum. Evidence-based practice is discussed within the context of emerging concerns over its application to non-medical interventions and suggestions are made for grounding fluency interventions by reference to empirically supported principles of change. A popular, evidence-based intervention for stuttering in young children (the Lidcombe program) is evaluated within the suggested parameters.

Educational objectives: After reading this article, the reader will be able to: (1) evaluate the status of evidence-based practice in fluency disorders; (2) list concerns about the impact of narrow interpretation of EBP on research and practice in the field of fluency disorders and other non-medical domains; (3) articulate the role of theory and basic research in selecting among and evaluating therapy approach options.

© 2005 Published by Elsevier Inc.

Keywords: Evidence-based practice; Treatment; Empirically-supported principles of change; Lidcombe

* Tel.: +1 301 405 4217.

E-mail addresses: nratner@hesp.umd.edu, nratner@aol.com (N. Bernstein Ratner).

In a recent issue of *JFD* (28, 3) (hereafter, 28/3), a number of authors (Bothe, 2003; Finn, 2003; Ingham, 2003; Onslow, 2003) proposed certain standards to which stuttering treatment should be held and further suggested that only a limited number of therapy programs currently meet such evidence-based practice (EBP) standards. In the same series of articles, some authors appeared to imply that treatment programs not meeting the specified standards, and the clinicians who administer them, may in fact be engaging in less than ethical clinical practice, since they volitionally forgo a small set of “validated” techniques for those seemingly supported by a lesser evidence base. In this space, I would like to further discuss such implications as well as related, seemingly provocative issues. I will also address larger issues in evaluating the degree to which EBP is currently fully “ready for prime time” implementation in the field of fluency disorders. In doing so, I will frame my comments in the form of a series of questions that I think we need to ask and consider answering before applying some of the extended principles of EBP to the field of stuttering intervention.

On the surface, EBP is a noble concept and goal. Indeed, it would seem nonsensical to argue for therapeutic practice that is not based on some body of evidence. I will not, therefore, position myself as saying that EBP is wrong. At the same time, certain embodiments and extensions of EBP seem less obviously of value and may in fact pose difficulties for researchers, clinicians and their patients.

In framing this article partially as a response to the authors in 28/3, I prefer to start with areas of agreement. I wholeheartedly concur with the obvious need for practitioners to document the rationale for their selection of therapy approaches. As such, I also most wholeheartedly agree with the general consensus of the authors in the issue that we need far more research into therapeutic efficacy in stuttering treatment. Thus, I agree with Ingham (2003) that researchers need to develop more interest in therapy trials, and that our funding agencies, particularly the American National Institutes of Health, need to invest in them more aggressively. It is possible to conjecture that the lack of funded research in therapy efficacy is at least in part due to the relative paucity of such applications when weighed against the bulk of submissions that propose basic research questions. Despite these strong areas of agreement, however, my affinity for some, if not many, of the arguments raised in the issue begins to wane considerably, because they raise a number of vexing questions. I address what I view to be the most important of these questions in the remainder of this article. Among the issues that I will consider are:

- 1) The nature and scope of “evidence” and its relationship to clinical practice;
- 2) The limitations that may be associated with the use of a single framework to implement EBP across medicine and the many health-related professions;
- 3) The role of different types of evidence in determining the value of specific therapy approaches in stuttering;
- 4) The role of theory in evaluating treatment approaches;
- 5) Potential barriers to the gathering of clinical evidence and its implementation by practitioners; and finally,
- 6) Some logical “next steps” that will be required if practitioners and researchers are to bridge the perceived gaps between evidence and practice in stuttering treatment.

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات