



Contents lists available at ScienceDirect

Journal of Behavior Therapy and Experimental Psychiatry

journal homepage: www.elsevier.com/locate/jbtep

How we walk affects what we remember: Gait modifications through biofeedback change negative affective memory bias



Johannes Michalak^{a, b, *}, Katharina Rohde^b, Nikolaus F. Troje^c

^a Witten Herdecke University, Department of Psychology and Psychotherapy, Alfred-Herrhausen-Straße 50, D-58448 Witten, Germany

^b University of Hildesheim, Institute of Psychology, Department of Clinical Psychology, Marienburger Platz 22, D-31131 Hildesheim, Germany

^c Queen's University Kingston, Department of Psychology, Biomotion Lab, 62 Arch Street, Kingston, Ontario K7L 3N6, Canada

ARTICLE INFO

Article history:

Received 26 February 2014

Received in revised form

22 July 2014

Accepted 4 September 2014

Available online 28 September 2014

Keywords:

Embodiment

Depression

Body

Memory

Biofeedback

ABSTRACT

Background and objects: Several studies have shown that physical exercise such as walking has effects on depression. These studies have focused on increasing intensity and amount of physical activity. In the present study, we investigated whether not only the intensity but also the style of physical activity affects depression related processes.

Method: Using an unobtrusive biofeedback technique, we manipulated participants (39 undergraduates) to change their walking patterns to either reflect the characteristics of depressed patients or a particularly happy walking style. The intensity of walking (i.e. walking speed) was held constant across condition. During walking, participants first encoded and later recalled a series of emotionally loaded terms. **Results:** The difference between recalled positive and recalled negative words was much lower in participants who adopted a depressed walking style as compared to participants who walked as if they were happy.

Limitations: The effects of gait manipulation were investigated in a non-clinical group of undergraduates. **Conclusions:** The observed change in memory bias supports the idea that beyond the intensity of walking the style of walking has effects on the vulnerability to depression.

© 2014 Elsevier Ltd. All rights reserved.

1. Introduction

A number of studies have shown that physical exercise has effects on mood and can be therapeutic for patients suffering from Major Depressive Disorders (MDD). In a meta-analysis by Reed and Ones (2006) including 158 studies it was shown that engaging in acute exercise (e.g. walking) increases positive affect. In addition, Mata et al. (2012) found that natural variations in physical activity had effects on mood in depressed individuals. In this 1-week experience sampling study depressed individuals expressed higher levels of positive affect when they were physically active than during sedentary periods. Moreover, studies investigating physical exercise as a treatment approach have documented that engaging in prescribed exercise significantly reduces levels of depressive symptoms in clinical samples (Conney et al., 2013). Remarkably, the effects of exercise on MDD seem to be comparable to effects of

antidepressant medication (Babyak et al., 2000; Hoffman et al., 2011).

Most of these physical treatment approaches to MDD posit that the central operating mechanism is the increased intensity of activity (Ekkekakis, 2003). It is proposed that the low activity level, which is characteristic for depressed individuals, should be increased to achieve the positive effects of exercises. However, in addition to a generally low level of activity, depressed individuals often show a number of further motoric characteristics. One central motoric characteristic of depressed individuals is a deviant gait pattern. A number of studies have shown a close relationship between walking style and depression (Bader, Bühler, Endrass, Klipstein, & Hell, 1999; Lemke, Wendorff, Mieth, Buhl, & Linnemann, 2000; Paleacu et al., 2007; Sloman, Berridge, Homatidis, Hunter, & Duck, 1982; Sloman, Pierrynowski, Berridge, Tupling, & Flowers, 1987). In a comprehensive analysis of gait characteristics in patients suffering from a current episode of major depression, for example, it has been demonstrated that patients not only showed reduced walking speed (i.e., activity level) but also smaller arm-swing amplitudes, smaller amplitude of vertical movements of the upper body, larger amplitudes of lateral body sway, and a more slumped and forward-leaning posture than

* Corresponding author. Witten Herdecke University, Department of Psychology and Psychotherapy, Alfred-Herrhausen-Straße 50, D-58448 Witten, Germany. Tel.: +49 (0) 2302 926 780.

E-mail address: johannes.michalak@uni-wh.de (J. Michalak).

healthy controls (Michalak et al., 2009). An analysis of speed dependent changes in gait parameters in non-clinical samples (Thorstensson, Nilsson, Carlson, & Zomlefer, 1984) makes it unlikely that gait characteristics of depressed patients are solely attributable to reduced walking speed (i.e., the intensity of walking).

A large body of basic research points to the fact that not only changes in general activity level but also more subtle motoric changes can have profound effects on emotional information processing. Under the term ‘embodiment’ numerous studies have shown a complex and reciprocal relationship between the bodily expression of emotion and the way in which emotional information is processed (Niedenthal, 2007; Niedenthal, Barsalou, Winkielman, Krauth-Gruber, & Ric, 2005).

The fact that emotional processes affect bodily states seems to be almost trivial. Obviously, emotions are expressed by the facial muscular system, but also in the static posture and dynamic activity of the whole body (Oberman, Winkielman, & Ramachandran, 2007; Pitcher, Garrido, Walsh, & Duchaine, 2008). More interesting are demonstrations of causal relations in the other direction. Diverse affective and cognitive processes are affected by manipulations of the muscular-skeletal system. For instance, in a seminal study Strack, Martin, and Stepper (1988) asked participants to hold a pen between their teeth, which forced them to contract the zygomaticus major muscle that is normally involved in the production of a smile. They found that the participants enjoyed cartoons more than those who were prevented from making zygomaticus contractions. Similarly, Stepper and Strack (1993) demonstrated that a slumped versus an upright sitting posture affected the way participants perceived positive feedback on a previously completed achievement test.

Bodily processes can also influence affective memory. Riskind (1983) showed that certain postures affected the ease with which participants were able to recall either pleasant or unpleasant life experience and Förster and Strack (1996) demonstrated that nodding or shaking the head influences a person's recall of positive and negative words.

Although basic research on embodiment effects on emotional information processing is flourishing, research on the effects of motoric system on processes relevant for psychopathological phenomena is largely lacking. In the current study, we manipulated participants' gait to adopt features reflecting the characteristics of happy or depressed walking behaviour using a biofeedback method. We then measured how many negative and positive words participants recalled from a previously presented list of items. Biased processing of self-referent material is one of the most intensively studied cognitive-affective correlates of depression. A large body of research has found that depression affects how information about oneself is processed, as evidenced by enhanced encoding and recall of negative self-referent material and by a reduced processing of positive information (Mathews & MacLeod, 2005; Matt, Vázquez, & Campbell, 1992; Williams, Watts, MacLeod, & Mathews, 1997). We expected to observe a memory bias such that manipulating the participants to walk in a more depressed way would shift recall towards relatively more negative words, and making them walk happier would bias them towards remembering relatively more positive words. Walking speed was held constant across conditions to rule out that mere differences in activity level are responsible for group differences in memory bias. We choose memory bias as our primary outcome variable because we expected that the relatively short time gait changes we aimed to induce would have effects on more subtle emotional information processes (i.e., memory bias), before effects on conscious experienced affective states could be detected. However, we also assessed the conscious affective state in order to investigate whether the

linkage between motor behavior and affective processes was mediated by the explicit experience of affective states or whether the sensorimotor system can affect these functions directly before effects on conscious affect become apparent.

2. Method

2.1. Participants

A total of 47 volunteers participated in this study. All were students from Queen's University who received either course credit for their participation or a monetary compensation of \$20. Data from eight of the participants were not usable due to technical problems and resulting data loss. The analysis was therefore based on data from 39 participants.

2.2. Apparatus

An optical motion capture system (17 cameras, 120 Hz sampling rate, Qualisys, Gothenburg, Sweden) was used to acquire in real time full-body movements exhibited by the participants while walking on a treadmill at a constant speed of 1.2 m/s (Troje, 2002, 2008). Data were analyzed within a 2 s temporal window according to where it scaled along a sad-happy continuum (see [Supplementary Material](#)). The continuum was parametrized in terms of a linear discriminant function which was based on the data of 14 depressive and 14 healthy participants obtained in the context of a previous study and described in more detail there (Michalak et al., 2009; Troje, 2008). An interactive animation visualizing the linear discriminant function can be viewed at <http://biomotionlab.ca/Demos/BMLdepression.html>. While walking on a treadmill the participants were facing a monitor mounted on a wall 4 m in front of them. On the monitor, we displayed a large gauge consisting of a horizontal scale and a vertical bar that could move along that scale. The gauge was calibrated such that the bar was located at the centre position if the participant's score was the same as during the walk exhibited at the beginning of the experiment (baseline condition, see below). From this null position the gauge deflected to the right if the participant's score indicated a happier walk and it deflected to the left if the participant's score indicated a more depressed walking pattern. In half of the participants, this relation was reversed. In either case, participants were instructed to try to walk such that the gauge would deflect to the right. The gain of the gauge was 10 cm/std (z-score relative to the distribution used to compute the discriminant function, Michalak, et al., 2009). Scores were obtained every 100 ms and the display of the gauge was updated at the same frequency. Scores were also stored in order to assess to which degree the feedback altered the participant's walk.

2.3. Instruments

To measure the effects of gait feedback on affective memory bias we adapted the self-referent encoding task (SRET, Ramel et al., 2007). In an initial encoding phase, the experimenter read out a list of 20 positive and 20 negative words (see [Supplementary Material](#)) in random order and the participants had to decide whether or not each word described them well. During a subsequent recall phase, which was applied later during the experiment and was not anticipated by the participant, they were then asked to recall as many words as possible from the entire list. Affective memory bias was measured as the difference between the number of recalled positive words and the number of recalled negative words. After participants had completed all other parts of the experiment and had stopped walking on the treadmill, they were subjected to a final questionnaire designed to assess their affective

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات