

A Preliminary Investigation of the Relationship Between Emotion Regulation Difficulties and Posttraumatic Stress Symptoms

Matthew T. Tull

Center for Addictions, Personality, and Emotion Research and the University of Maryland

Heidi M. Barrett

University of Massachusetts Boston

Elaine S. McMillan

University of Maine

Lizabeth Roemer

University of Massachusetts Boston

This study examined the relationship between posttraumatic stress (PTS) symptoms and particular aspects of emotion regulation difficulties among trauma-exposed individuals. Participants were an ethnically diverse sample of 108 undergraduates from an urban university. PTS symptom severity was found to be associated with lack of emotional acceptance, difficulty engaging in goal-directed behavior when upset, impulse-control difficulties, limited access to effective emotion regulation strategies, and lack of emotional clarity. Further, overall difficulties in emotion regulation were associated with PTS symptom severity, controlling for negative affect. Finally, individuals exhibiting PTS symptoms indicative of a PTSD diagnosis reported greater difficulties with emotion regulation than those reporting PTS symptoms at a subthreshold level. The implications of these findings for research and treatment are discussed.

EXPOSURE TO A POTENTIALLY traumatic event is defined in part by the experience of an intense emotional response (i.e., fear, helplessness, and horror; [American Psychiatric Association \[APA\], 1994](#)), and in the immediate aftermath of the event, it is common for individuals to experience heightened emotional and physiological responses to frequent reminders of the event (e.g., [Rothbaum, Foa, Riggs, Murdock, & Walsh, 1992](#); [Shalev et al., 1998](#)). Over time, these responses diminish for many, if not most, survivors ([Kessler, Sonnega, Bromet, Hughes, & Nelson, 1995](#); [Rothbaum et al., 1992](#); [Shalev et al., 1998](#)), whereas others go on to develop chronic symptoms of reexperiencing, avoidance, and hyperarousal ([APA, 1994](#)). A substantial body of research has attempted to identify factors that may be associated with the persistence of these posttraumatic stress (PTS) symptoms (for a review, see [Brewin, Andrews, & Valentine, 2000](#)); however, little attention has been paid to the role of difficulties in the ways in which individuals respond to or regulate their intense emotions.

Theorists have highlighted the potential importance of emotion regulation difficulties in PTS and PTSD ([Cloitre, 1998](#); [Cloitre, Koenen, Cohen, & Han, 2002](#); [van der Kolk, 1996](#)). Further, findings of an association between posttraumatic stress disorder (PTSD) and more intense emotional (subjective and psychophysiological) responding

Preparation of this manuscript was supported in part by training grant MH20048 from the National Institute of Mental Health of the National Institutes of Health to the first author.

Portions of this study were presented at the 39th annual meeting of the Association for Behavioral and Cognitive Therapies in Washington, DC (2005).

Address correspondence to Mathew T. Tull, Department of Psychology, University of Maryland, College Park, MD 20742, USA; email: MTull@psyc.umd.edu.

0005-7894/07/0303-0313\$1.00/0

© 2007 Association for Behavioral and Cognitive Therapies. Published by Elsevier Ltd. All rights reserved.

to emotionally evocative stimuli (Litz, Orsillo, Kaloupek, & Weathers, 2000; Orsillo, Batten, Plumb, Luterek, & Roessner, 2004; McDonagh-Coyle et al., 2001; Veazey, Blanchard, Hickling, & Buckley, 2004) indicate that PTS symptoms might be associated with heightened emotional responses, which would then require greater regulation efforts (Mennin, 2005). An inability to effectively down-regulate this emotional arousal or difficulties in the awareness and differentiation of emotional states may contribute to the perception of emotions as uncontrollable and unpredictable—two factors that have been found to influence the likelihood of fear acquisition (Bouton, Mineka, & Barlow, 2001). Consequently, individuals may learn to fear (and subsequently avoid) situations where certain emotions may be elicited, thus preventing functional exposure to trauma-relevant cues (Foa & Kozak, 1986). In addition, difficulties in emotion regulation may indirectly influence the maintenance of PTS symptoms (e.g., avoidance) through their negative effect on interpersonal relationships and an individual's overall functional capacity (see Cloitre et al., 2002). Thus, difficulties in emotion regulation abilities might serve to maintain post-traumatic symptoms in multiple ways. However, the particular difficulties in emotion regulation (using a comprehensive definition) that may be associated with posttraumatic responding have yet to be extensively explored (Hunt and Evans, 2004). Examination of the relationship between emotion regulation difficulties and PTS symptoms is clinically important and may enhance our understanding of the maintenance of PTS symptoms.

Our conceptualization of emotion regulation emphasizes the functional nature of emotional responses, with regulation referring to an ability to act effectively in the context of emotionally salient events. Gratz and Roemer (2004), consistent with Thompson (1994), broadly define emotion regulation as the ability to monitor, evaluate, and modulate emotional reactions, especially in the context of goal-directed behavior. As such, they propose that effective emotion regulation involves several domains, including: (a) the awareness and understanding of emotions, (b) the acceptance of emotions, (c) the ability to control impulsive behaviors and engage in goal-congruent behaviors in the context of distressing emotional experiences, and (d) access to and flexible use of situationally appropriate emotion regulation strategies. Empirical evidence supports the distinct but related nature of these domains (Gratz & Roemer, 2004).

Inherent in this definition of emotion regulation is the distinction between the modulation and

avoidance of an emotion. Modulating an emotion involves altering the intensity or duration of an emotion as opposed to attempting to avoid or extinguish the emotion altogether (which is commonly associated with psychopathology and impairment rather than emotional health; see Hayes, Luoma, Bond, Masuda, & Lillis, 2006, for a review). Monitoring and evaluating emotional reactions are also important regulatory components; understanding and accepting one's emotions are seen as crucial components of emotion regulation, as nonacceptance or lack of understanding may prolong emotional distress by adding negative secondary emotions to the primary negative emotion (e.g., feeling shameful about having the experience of anxiety or fear; Gratz & Roemer, 2004; Greenberg & Safran, 1987; Hayes, Strosahl, & Wilson, 1999). For example, Salters-Pedneault, Gentes, and Roemer (2007) found that self-reported trait levels of fear of depression, anxiety, and positive emotions predicted increased subjective and physiological reactivity to a trauma-related, emotionally evocative stimulus. Further, given that many individuals experience their emotions as preventing them from carrying out necessary activities, the emphasis on behavioral responding in the face of emotional experience (e.g., inhibiting impulses and pursuing goals when upset) is especially important. An individual's behavior and ability to pursue self-care, pleasure, and mastery activities is an important indicator of an individual's ability to regulate emotions (Gross & Munoz, 1995).

Trauma-exposed individuals who are faced with memories of emotionally evocative events may struggle with each aspect of emotion regulation. Evidence suggests an association between PTSD and elevations in alexithymia, which is defined by difficulty identifying feelings, as well as a limited ability to distinguish between feelings and bodily sensations of emotional arousal (Taylor, Bagby, & Parker, 1991). Several studies have revealed significant positive associations between the severity of PTSD, traumatic exposure, and alexithymia (e.g., Fukunishi, Sasaki, Yasunori, Masanori, & Masaki, 1996; Yehuda et al., 1997). Further, Monson, Price, Rodriguez, Ripley, and Warner (2004) found that difficulty describing emotions (combined with the avoidance of internal emotional experiences) was associated with greater PTS symptom severity, especially for reexperiencing symptoms. These findings suggest that poor understanding of emotions may be associated with PTS.

Although no studies have directly tested the relationship between the behavioral aspect of emotion regulation difficulties (i.e., an inability to

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات