



The effects of childhood abuse on symptom complexity in a clinical sample: Mediating effects of emotion regulation difficulties[☆]



Ji Young Choi^a, Young Min Choi^a, Min Sook Gim^a, Jun Hyun Park^a,
Soo Hyun Park^{b,*}

^a Department of Psychiatry, Sanggye Paik Hospital, Inje University, 1342, Dongilro, Nowon-gu, Seoul, Republic of Korea

^b Department of Psychology, Yonsei University, Shinchon-dong, Seodaemun-gu, Seoul, Republic of Korea

ARTICLE INFO

Article history:

Received 20 January 2014

Received in revised form 21 April 2014

Accepted 23 April 2014

Available online 17 May 2014

Keywords:

Childhood abuse

Symptom complexity

Emotion regulation difficulty

ABSTRACT

The purpose of the present study was to first examine whether childhood abuse predicts symptom complexity, as indicated by the number of clinically elevated scales on the MMPI-2 in an adult clinical sample. Secondly, we investigated whether emotion regulation difficulties mediated the relationship between childhood abuse and symptom complexity. A total of 162 adult outpatients not presenting with psychotic symptoms completed the Korean Childhood Trauma Questionnaire (K-CTQ), Life Events Checklist (LEC), Difficulties in Emotion Regulation Scale (DERS), and Korean Minnesota Multiphasic Personality Inventory-2 (MMPI-2). Partial correlation analysis results indicated that after controlling for the presence of adulthood trauma, childhood abuse was associated with more symptom complexity, or more clinically elevated scales on the MMPI-2. Furthermore, structural equation modeling results showed that emotion regulation difficulties partially mediated the relationship between childhood abuse and symptom complexity. These findings indicate that individuals who had experienced childhood abuse evidence simultaneous presentation of diverse clinical symptoms.

© 2014 Elsevier Ltd. All rights reserved.

Introduction

It has been widely recognized that childhood abuse serves as a significant risk factor for maladjustment and psychopathology in adulthood (Briere & Rickards, 2007; Cicchetti, Ackerman, & Izard, 1995). Studies focusing on community and clinical samples consistently report a significant correlation among childhood emotional, physical, and sexual abuse and impairment in diverse functional domains, as well as indices of psychopathology such as posttraumatic stress, depression, substance abuse, and suicidal ideation in adulthood (Briere & Runtz, 1988; Fergusson, Boden, & Horwood, 2008; Min, Farkas, Minnes, & Singer, 2007; Rowan, Foy, Rodriguez, & Ryan, 1994; Schaaf & McCanne, 1998).

Interpersonal trauma such as childhood abuse is associated with symptom sequelae which are different from typical posttraumatic stress disorder (PTSD), and thus, such a clinical presentation has been posited to be a unique variation of PTSD. Herman (1992) and Pelcovitz and colleagues (1997) had proposed an independent diagnosis of Complex PTSD or

[☆] This work was supported by the 2013 Inje University research grant. The funding source had no involvement in the conduct of research and preparation of the article. Poster presentation of a portion of this research was made at the 2014 American Psychological Association Annual Convention.

* Corresponding author.

Disorders of Extreme Stress Not Otherwise Specified (DESNOS) manifested as alterations in six areas of functioning such as regulation of affect and impulses, attention and consciousness, self-perception, somatization, relations with others, and systems of meaning. Such propositions attempted to explain the consequences of cumulative childhood trauma as a unique pathological entity.

If Complex PTSD or DESNOS focused on the psychopathological presentation as an independent specific diagnostic entity, the concept of symptom complexity can be construed as a more comprehensive construct encompassing more diverse symptoms that are presented simultaneously. Indeed, several researchers have focused on the complexity in symptomatology as a unique negative consequence of childhood trauma. Symptom complexity has been defined as more different kinds of symptoms that occur simultaneously, or *multisymptom clinical presentations* (Briere, Kaltman, & Green, 2008). Thus, symptom complexity refers to a complex symptom presentation that includes not only difficulties related to mood or social withdrawal, reflecting disturbances predominantly in affective and interpersonal domains, but also symptoms such as destructive behavior and anger (Cloitre et al., 2009). In this vein, symptom complexity does not reflect the severity of clinical symptomatology per se but presentation of multiple symptoms across diverse domains of psychopathology.

For instance, greater number of cumulative childhood trauma was positively associated with clinical levels of symptomatology in 10 different symptom subdomains such as anxiety arousal, depression, sensitivity, and re-experiencing on the Trauma Symptom Inventory (Briere et al., 2008). This finding suggests that childhood trauma is related to symptom complexity in adulthood. Cloitre and colleagues (2009) examined whether more types of childhood trauma experienced (e.g., sexual abuse, physical abuse, neglect, emotional abuse, separation from mother) is associated with symptom complexity in a sample of women seeking treatment for abuse. Symptom complexity in this study was assessed through the number of clinically elevated levels of symptomatology in six symptom types across three domains such as symptoms of posttraumatic stress, emotion dysregulation, and problems in interpersonal relationship.

Interest in symptom complexity shares the context of Complex PTSD or DESNOS in that repeated trauma that persists from the early phase of development leads to complex symptomatology secondary to not only disruption in self-concept or trust but also impairment in basic emotion regulation capacity. Emotion regulation has been defined in diverse ways, but in general, it has been regarded as both an internal and external process wherein an individual monitors, evaluates, corrects and modulates emotional responses, in particular emotional responses characterized by intense and transient nature for the purpose of attaining a goal (Thompson, 1994). Extant studies have consistently demonstrated that difficulties in emotion regulation is associated with diverse psychological problems such as depression, anxiety, and posttraumatic stress symptoms (Eftekhari, Zoellner, & Vigil, 2009; McLaughlin, Hatzenbuehler, Mennin, & Nolen-Hoeksema, 2011; Price, Monson, Callahan, & Rodriguez, 2006).

It has been demonstrated that higher-order cortical structures, particularly the prefrontal cortex and its subunits, mediate attempts to regulate emotions, structures implicated in studies examining fear and anxiety (Quirk & Beer, 2006). In the case of trauma-related stress response, overwhelming anxiety or intense negative affective experience not only triggers PTSD symptoms such as anxiety-arousal response but also general emotion regulation difficulties such as reactivity and arousal regulation problems (Ford, Wasser, & Connor, 2011). In other words, trauma disturbs the anxiety-arousal response system and in this vein, the re-experiencing and hyperarousal responses characteristic of PTSD may also be considered as a form of emotion regulation difficulty that is associated with diverse psychological problems and general functional impairment (Tull, Barrett, McMillan, & Roemer, 2007). In particular, abuse experience in childhood when emotion regulation skills have not been adequately acquired or developed may magnify changes in reactivity and ultimately interfere with children's achievement of adequate self-soothing and other emotion regulation strategies (Cloitre, Miranda, Stovall-McClough, & Han, 2005; Kim & Cicchetti, 2009; Maughan & Cicchetti, 2002). This notion has been suggested as one reason why there exists a greater likelihood of survivors of childhood trauma to manifest more complex clinical symptomatology in adulthood in addition to the characteristic emotion regulation difficulties.

Recent studies have looked more particularly into the role of emotion regulation as both a direct negative consequence of childhood abuse and also as a mediator in the relationship between childhood abuse and subsequent psychological problems. Abused children evidenced more externalizing problems such as aggressive behavior as well as more internalizing problems such as depression and anxiety and such results were mediated by impairments in emotion regulation (Schwartz & Proctor, 2000; Shields & Cicchetti, 2001). Such impairment in emotion regulation also mediated the relationship between cumulative trauma and behavior problems (Choi and Oh, 2014). Childhood interpersonal trauma significantly predicted emotion regulation difficulties and such difficulties mediated the relationship between childhood interpersonal trauma and interpersonal functioning and many adjustment-related problems (Briere & Rickards, 2007; Cloitre et al., 2005).

In summary, extant research suggests the childhood trauma such as abuse results in significant negative consequences and that impairment in emotion regulation is both a negative consequence and an important mechanism connecting the experience of childhood trauma with subsequent difficulties. Furthermore, one of the unique features of such negative consequence of childhood trauma is the manifestation of symptom complexity. However, extant research has mostly focused on complex PTSD as reflective of symptom complexity. This precludes the possibility that symptom presentation related to childhood abuse can be simultaneously manifested across diverse symptom domains in the actual clinical setting. As such, using a tool such as the Minnesota Multiphasic Personality Inventory-2 (MMPI-2), which can measure not only symptoms of affective or relational dysfunction but more comprehensive psychiatric symptoms and personality-related issues, may assist in evaluating symptom complexity. The MMPI-2 holds the advantage as a well standardized and validated tool that is applicable to diverse clinical samples and capable of assessing diverse clinical symptoms. One study utilized the MMPI

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات