



Review

Biological and psychological markers of stress in humans: Focus on the Trier Social Stress Test



Andrew P. Allen^{a,b}, Paul J. Kennedy^{a,b}, John F. Cryan^{a,c},
Timothy G. Dinan^{a,b}, Gerard Clarke^{a,b,*}

^a Alimentary Pharmabiotic Centre, University College Cork, Cork, Ireland

^b Department of Psychiatry, University College Cork, Cork, Ireland

^c Department of Anatomy and Neuroscience, University College Cork, Cork, Ireland

ARTICLE INFO

Article history:

Received 1 August 2013

Received in revised form 1 November 2013

Accepted 6 November 2013

Keywords:

Stress

Cortisol

ACTH

Serotonin

HPA axis

Sympathetic-adrenal-medullary system

Brain–gut axis

Heart rate

Immune system

Genetics

ABSTRACT

Validated biological and psychological markers of acute stress in humans are an important tool in translational research. The Trier Social Stress Test (TSST), involving public interview and mental arithmetic performance, is among the most popular methods of inducing acute stress in experimental settings, and reliably increases hypothalamic–pituitary–adrenal axis activation. However, although much research has focused on HPA axis activity, the TSST also affects the sympathetic-adrenal-medullary system, the immune system, cardiovascular outputs, gastric function and cognition. We critically assess the utility of different biological and psychological markers, with guidance for future research, and discuss factors which can moderate TSST effects. We outline the effects of the TSST in stress-related disorders, and if these responses can be abrogated by pharmacological and psychological treatments. Modified TSST protocols are discussed, and the TSST is compared to alternative methods of inducing acute stress. Our analysis suggests that multiple readouts are necessary to derive maximum information; this strategy will enhance our understanding of the psychobiology of stress and provide the means to assess novel therapeutic agents.

© 2013 Elsevier Ltd. All rights reserved.

Contents

| | |
|--|-----|
| 1. Introduction | 95 |
| 1.1. General introduction | 95 |
| 1.2. The Trier Social Stress Test | 96 |
| 2. Biomarkers and psychological effects of the TSST | 96 |
| 2.1. Biomarkers | 96 |
| 2.1.1. Hypothalamic–pituitary–adrenal (HPA) axis effects | 96 |
| 2.1.2. Immune system effects | 97 |
| 2.1.3. Sympathetic-adrenal-medullary system effects | 98 |
| 2.1.4. Cardiovascular effects | 99 |
| 2.1.5. Brain–gut axis effects | 99 |
| 2.1.6. Biomarkers: summary | 99 |
| 2.2. Psychological effects | 99 |
| 2.2.1. Subjective effects | 99 |
| 2.2.2. Cognitive effects | 99 |
| 2.2.3. Neurobiological effects | 100 |
| 2.2.4. Psychological effects: summary | 100 |
| 3. Procedural considerations | 100 |
| 3.1. Biomarkers | 100 |
| 3.1.1. HPA axis measures | 100 |

* Corresponding author at: Department of Psychiatry, 1.15 Biosciences Institute, University College Cork, Cork, Ireland. Tel.: +353 21 4901408.
E-mail address: G.Clarke@ucc.ie (G. Clarke).

| | | |
|--------|---|-----|
| 3.1.2. | Immune system measures | 101 |
| 3.1.3. | Sympathetic-adrenal-medullary system measures | 101 |
| 3.1.4. | Cardiovascular measures | 101 |
| 3.1.5. | Brain–gut axis measures | 102 |
| 3.1.6. | Procedural considerations in biomarkers: summary | 102 |
| 3.2. | Psychological effects | 102 |
| 3.2.1. | Subjective measures | 102 |
| 3.2.2. | Cognitive measures | 102 |
| 3.2.3. | Procedural considerations in psychological effects: summary | 103 |
| 4. | Factors moderating the effects of the TSST | 103 |
| 4.1. | Gender and age | 103 |
| 4.1.1. | Age and gender differences | 103 |
| 4.1.2. | Sex hormones and menstrual cycle | 103 |
| 4.2. | Genetics | 103 |
| 4.3. | Personality and individual differences | 104 |
| 4.3.1. | Personality factors | 104 |
| 4.3.2. | Stressor appraisal | 104 |
| 4.4. | Social factors | 104 |
| 4.4.1. | The interview panel | 104 |
| 4.4.2. | Social support and social status | 104 |
| 4.4.3. | Cultural factors | 105 |
| 4.5. | Factors moderating the effects of the TSST: summary | 105 |
| 5. | TSST response and stress-related psychiatric disorders | 105 |
| 5.1. | Depression | 105 |
| 5.2. | Anxiety | 105 |
| 5.3. | Other stress-related psychiatric disorders | 106 |
| 5.4. | Factors moderating TSST response in stress-related disorders | 106 |
| 5.5. | Treatment for stress-related psychiatric disorders | 106 |
| 5.5.1. | Psychological treatment for stress-related psychiatric disorders | 106 |
| 5.5.2. | Pharmacological treatment for stress-related psychiatric disorders | 107 |
| 5.5.3. | Dietary interventions for stress | 107 |
| 5.6. | TSST and stress-related psychiatric disorders: summary | 107 |
| 6. | Modified versions of the TSST | 107 |
| 6.1. | Repeated measures | 107 |
| 6.2. | Control TSST | 111 |
| 6.3. | Group assessment | 111 |
| 6.4. | Children and adolescents | 111 |
| 6.5. | Virtual/imagined TSST committee | 112 |
| 6.6. | Hybrid versions of the TSST | 112 |
| 6.7. | Modified versions of the TSST: summary | 112 |
| 7. | Discussion | 112 |
| 7.1. | Comparison with other methods of inducing stress and measuring HPA axis activity | 112 |
| 7.1.1. | Pharmacological stimulation of the HPA axis | 112 |
| 7.1.2. | The cold pressor task | 113 |
| 7.1.3. | The CO ₂ challenge test | 113 |
| 7.1.4. | The cortisol awakening response | 113 |
| 7.1.5. | Cognitive tasks | 113 |
| 7.1.6. | Noise | 113 |
| 7.1.7. | Comparison with other methods of inducing stress and measuring HPA axis activity: summary | 114 |
| 7.2. | Future research | 114 |
| 7.3. | Summary and conclusions | 115 |
| | Acknowledgements | 117 |
| | References | 117 |

1. Introduction

1.1. General introduction

Exposure to a situation perceived as threatening or excessively demanding (a stressor) leads to the release of chemicals which can help cope with the stressor (e.g. Joels and Baram, 2009; Schwabe et al., 2012). Acute stress can alter numerous biological functions, such as the hypothalamic-pituitary-adrenal (HPA) axis (Foley and Kirschbaum, 2010), the immune system (Steptoe et al., 2007), the autonomic nervous system (Xhyheri et al., 2012), and the enteric nervous system (Ziegler, 2012). Psychologically, acute stress is a subjectively negative experience (but see Jamieson et al., 2013), and can have positive as well as negative effects on cognition (Lupien

et al., 2007; Starcke and Brand, 2012). Although acute stress is important for responding to threatening situations, chronic stress is damaging to health (McEwen, 2007). Over time, stressful life events can induce chronic stimulation of the HPA axis, leading to depression (Checkley, 1992; Juruena et al., 2011), and poorer prognosis for cancer and heart disease (Maddock and Pariante, 2001). Early-life stress is associated with disorders such as depression (Batten et al., 2004; Dinan, 2005), post-traumatic stress disorder (Koenen et al., 2007), and irritable bowel syndrome (Mayer et al., 2001), and chronic work stress can exacerbate anxiety and depression (Smith, 2000). Moreover, chronic stress will result in changes in subsequent reactivity to an acute stressor (e.g. Chatkoff et al., 2010; Low et al., 2009; Roth et al., 2012). Given the contrast between acute and chronic stress, it is of interest to examine the effects

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات