



Hair cortisol: A new tool for evaluating stress in programs of stress management



Silvia Iglesias^a, Darío Jacobsen^b, Diego Gonzalez^b, Sergio Azzara^c, Esteban M. Repetto^b, Juan Jamardo^b, Sabrina Garín Gómez^b, Viviana Mesch^b, Gabriela Berg^b, Bibiana Fabre^{b,*}

^a Faculty of Pharmacy and Biochemistry, Universidad de Buenos Aires, Argentina

^b Clinical Biochemistry Department, INFIBIOC, Faculty of Pharmacy and Biochemistry, Universidad de Buenos Aires, Argentina

^c Faculty of Psychology, Universidad de Buenos Aires, Argentina

ARTICLE INFO

Article history:

Received 21 May 2015

Received in revised form 14 September 2015

Accepted 6 October 2015

Available online xxxx

Keywords:

Hair cortisol

Salivary cortisol

Stress management

ABSTRACT

Aims: Longitudinal and experimental studies have shown that chronic stress contributes to the onset and progression of different diseases. Although it is not possible to eliminate stress completely, people can learn to manage it by participating in different kinds of stress management interventions. This study examined the effectiveness of stress management interventions on neuroendocrine responses in stressed students and health professionals, by measuring hair cortisol in comparison to salivary cortisol.

Main methods: Salivary and hair cortisol measurements were performed in 37 subjects (31 women, 6 men; mean age 34.0 ± 10.6) who attended to a Coping Stress and Quality of Care Program at the University of Buenos Aires. Cortisol was measured at the beginning and at the end of the program. The State-Trait Anxiety Inventory STAI was used to evaluate state and trait anxiety.

Key findings: In subjects who completed the program, no differences were observed in salivary cortisol levels between the first and the last session. However, in these subjects, hair cortisol obtained in the last session was significantly lower than hair cortisol in the first session.

Significance: Hair cortisol appears to be a better biomarker than salivary cortisol for evaluation of the effectiveness of a stress reduction program and it seems to be a better indicator of stress system dysregulation as well.

© 2015 Elsevier Inc. All rights reserved.

1. Introduction

Longitudinal and experimental studies have shown that chronic stress contributes to the onset and progression of different diseases [7,9,23,24,34]. The relationship between stress and the onset or prognosis of different diseases, such as the metabolic syndrome (MS), is quite established [7]. In fact, in cardiovascular disease (CVD), risk associated with psychosocial stress factors is similar to traditional CVD risk factors [43].

Although it is not possible to eliminate stress completely, people can learn to manage it by participating in different kinds of stress management interventions [1,13,40]. These interventions include cognitive behavioral therapies [8] and/or special practices such as mind–body techniques [19,21]. In experimental studies, the effectiveness of these interventions at workplace settings has been determined using variables derived from psychological tests (e.g., stress, anxiety, or depression tests) and, to a lesser extent, using physiological measurements (e.g., blood pressure, heart rate, salivary cortisol, galvanic skin response) [27].

Stressful stimuli can activate neural and neuroendocrine pathways. Glucocorticoids are commonly used as biomarkers of stress [29,30]. A

blunted cortisol response is associated with negative health outcomes [26] and it was found in patients with panic disorder under psychosocial stress [25], as well as in people with depressive and anxiety disorders [38], or diabetes mellitus [3] and tinnitus [11]. Among the different samples used for the measurement of cortisol, morning saliva provides a measurement at a single point in time and considering its major physiological daily fluctuations, it does not reflect the stress response for extended periods of time. Recently, the use of hair cortisol measurement demonstrated that it provides a retrospective index of integrated cortisol secretion over periods of several months and it was also described as a potential biomarker of chronic stress [28,33].

The aim of this study was to examine the effectiveness of stress management interventions on neuroendocrine responses in students and health professionals through the measurement of hair cortisol in comparison to salivary cortisol.

2. Methods

2.1. Participants

In this study, 83 subjects (71 women, 12 men; mean age 35.7 ± 12.2) were voluntarily enrolled through a recruiting e-mail sent to the

* Corresponding author at: Junin 956 CABA, Argentina.
E-mail address: bfabre@fyb.uba.ar (B. Fabre).

databases of Schools of Medicine, Biochemistry and Psychology, University of Buenos Aires (UBA) to attend to a Coping Stress and Quality of Care Program at this university. Health professionals and students were included in the study. Subjects were excluded if they refused to sign the informed consent form or if they presented serious underlying medical or psychological conditions that could interfere with the results of the study (such as Cushing's syndrome, depression or generalized anxiety disorder). Additional exclusion criteria were dyed hair and/or shorter than 3 cm hair length. The training program was considered completed with at least 70% of attendance. Only 67 subjects (58 women, 9 men; mean age 36.3 ± 12.3) attended the first meeting and 37 completed the program. Salivary and hair cortisol measurements were performed in 37 subjects (31 women, 6 men; mean age 34.0 ± 10.6). One of the limitations of this study could be the small sample size and the fact that it was carried out with a majority of female participants another, taking into account the fact that the associations of psychosocial stress with endocrine activities may differ between genders [39].

The participants did not receive any kind of compensation for participating in the study and all of them gave written prior informed consent. The study was approved in advance by the Ethics Committee of the hospital and was performed following the Helsinki Declaration for medical studies in humans.

2.2. Program

The program intervention consisted in stress reduction lessons. The group received 90–120 minute training sessions during 10 weeks. All participants received a training manual containing a program summary. The aim of the program was to teach a variety of skills that each subject could integrate into his or her life on a regular basis and they were encouraged to practice them outside the sessions. These sessions were divided into a day topic introduction (40–50 min per session), formal practice of deep breath, relaxation, meditation-guided imagery exercises (20–30 min per session), and a final reflective group discussion (30–40 min per session). The program included the following topics: stress, stress symptoms, physiology of stress and relaxation response, mind/body connection, diaphragmatic and deep breathing, control vs. stress, cognitive restructuring, personal problem-solving and time management [13].

2.2.1. Anxiety levels

The State-Trait Anxiety Inventory STAI [17,32] was used to evaluate state and trait anxiety. This test differentiates between transient state anxiety (STAI-S) and more stable trait anxiety (STAI-T), with respondents reporting how they feel at the time the test is performed and in general in their daily life. Each subscale comprises 20 items, with total scores ranging from 20 to 80. Usually levels between 20 and 40 are associated with moderate anxiety [5,17].

2.3. Measurements

2.3.1. Biochemical determinations

Salivary and hair cortisol were measured at the beginning (pre sample, first session) and at the end (post sample, last session) of the program.

2.3.2. Salivary cortisol

Saliva samples were obtained by spontaneous salivation immediately after awaking, 30 min after awaking and before bedtime (at 11 pm) in order to measure salivary cortisol, which was determined by electrochemiluminescence (Cobas e411 autoanalyzer, Roche Diagnostics, Mannheim, Germany) The results were expressed in nmol/L. Saliva sampling compliance was evaluated by participants' reports.

2.3.3. Hair cortisol

Hair samples were obtained from the posterior vertex, hair was cut in the area closer to the scalp. Once the samples were obtained, three centimeters were measured from the root segment adjacent to the cutting. Then, each sample was weighed and a minimum of 20 mg of hair was needed for a proper extraction. Cortisol was extracted by shaking and subsequently overnight incubation, using methanol as extraction solvent. An aliquot of the extract was taken out and the solvent evaporated before being reconstituted and processed by an automated method (Cobas e411 autoanalyzer, Roche Diagnostics, Mannheim, Germany). The results were expressed in pg/mg.

Initial hair samples were collected at the beginning of the first session of the program, then, the participants spent three months in the program, and end hair samples were collected at the end of the last session. So, there was a period of three months between pre- and post-program sampling. Hair grows on average about one centimeter per month, that is why measuring the cortisol in the first three centimeters of hair closer to the scalp would represent cortisol exposure during the past three months.

Roche Cobas e-411 Cortisol assay is standardized with Cortisol Enzymun-Test, which is standardized through isotopic dilution-mass spectrometry (ID-MS). The analytical sensitivity (limit of detection) is 0.5 nmol/L and it shows the following cross-reactivities: corticosterone 5.8%; cortisol-21-sulfate 0.04%; cortisone 0.30%; 11-deoxycorticosterone 0.69%; 11-deoxycortisol 4.1%; dexametasone 0.08%; 17- α -hydroxyprogesterone 1.50%; prednisone 0.28%; progesterone 0.35%. The quality control used was BIO-RAD Liphocheck Immunoassay Plus Control, Lot number 48,280.

2.3.4. Statistical methods

We first tested the distribution of variables using normality tests (kurtosis and skewness), and then performed transformations in order to normalize the data. Pearson correlations were computed between dependent and independent variables and between dependent variables and potential confounders. T-test for independent samples was performed between participants that completed the program and those that did not. Wilcoxon signed-rank test was used to analyze differences in pre- and post-program variables. A p-value of less than 0.05 was considered as statistically significant. The Statistical Package for Social Sciences (SPSS: version 17.0) was used for data analysis.

3. Results

3.1. Comparison of groups at baseline

Socio-demographic and anthropometric characteristics of the subjects who attended the first meeting are shown in Table 1. Regarding age and gender distribution, no differences were observed between the subjects who completed the program and those who did not.

Although psychophysiological variables showed lower values in the subjects who completed the program, these differences were not statistically significant (Table 1). Among the subjects who attended the first meeting, 70% reported the need to incorporate tools to better manage stressful situations as they considered themselves as stressed.

3.2. Effects of stress reduction programs

Pre- and post-program sample values and the effects of stress reduction programs in participants who completed the program and those that did not, are shown in Table 2. In those subjects who completed the program, no differences were observed in salivary cortisol at any time (basal, 30 min, 11 pm) between first and last session, (Fig. 1). However, hair cortisol from the last session was significantly lower than hair cortisol from the first session in these subjects (Table 2, Fig. 2). No correlations were found between any other variable.

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات