



Developing a soil quality index to compare soil fitness for agricultural use under different managements in the Mediterranean environment

E. Armenise^{a,*}, M.A. Redmile-Gordon^b, A.M. Stellacci^c, A. Ciccacese^a, P. Rubino^a

^a Department of Crop Science, University of Bari, Bari, Italy

^b Soil Science Department, Rothamsted Research, Harpenden, UK

^c Consiglio per la ricerca e la sperimentazione in agricoltura, Unità di ricerca per i sistemi colturali degli ambienti caldo-aridi, Bari, Italy

ARTICLE INFO

Article history:

Received 1 July 2012

Received in revised form 21 February 2013

Accepted 28 February 2013

Keywords:

Soil quality

Soil quality indicators

Water extractable organic C

Water extractable N

S index

Long-term field trial

ABSTRACT

Due to increasing land-use pressures, soil-quality assessment is in growing demand, thus a standard set of procedures to assign a soil quality index (SQI) would be beneficial. In this study, the effectiveness of different managements in upholding soil quality (for crop productivity) was compared using soil quality indicators incorporated systematically to determine a SQI. The managements included three cropping sequences (wheat based three-year crop rotation, wheat + dry bean double-cropping and continuous wheat), two stubble managements (burning vs. incorporation), and three NPK-rates (nil, intermediate and optimal). Soil physical and chemical parameters were measured, screened through principal component analysis (PCA), normalised, and then integrated into a weighted-additive SQI. Crop sequence significantly affected soil pH and had a measurable effect on plant available P, with lower pH and higher P availability for the legume-based annual rotation. Stubble incorporation enhanced the labile N pool in spring, while ashes generated from burning of residues increased the level of exchangeable cations. Changes in total N were not detectable for any management. The SQI indicated that soil quality was most affected by NPK rate. The correlation between SQI and yield was not statistically significant, suggesting that other soil quality indicators, not measured in this study, were more influential upon yield at the experimental site in 2006. The SQI obtained using the method described herein was able to synthesise the complex information contained in large multivariate data-sets, and therefore would be useful for application at regional and national scales.

© 2013 Elsevier B.V. All rights reserved.

1. Introduction

Since the early 1990s the concept of soil quality has received much attention. Great efforts were first made to define soil quality, then to turn this vague concept into something tangible. The soil quality index (SQI) comes directly from the need of a science-based tool to measure soil quality.

Soil quality has been defined as the capacity of a soil to function, within land use and ecosystem boundaries, to sustain biological productivity, maintain environmental quality, and promote plant, animal, and human health (Doran and Parkin, 1994). Unlike air or water, soil is not directly consumed by humans and animals, and therefore soil quality standards are more complicated to identify (Doran et al., 1996). It is thus not surprising that while EU legislation regarding air and water quality is well established, a proposal for a Soil Framework Directive only arrived in 2006

(EC, 2006) and is still under adjustment. It is clear that the standardisation of a method for assessing soil quality is not simply of scientific interest, but a requirement to empower much needed decision-making processes. Soil quality assessment by means of indices has been successfully adopted both at regional scale and on-farm level (Glover et al., 2000; Karlen et al., 1994, 2008; Mastro et al., 2008; Wienhold et al., 2006). We believe that this kind of approach may be useful in supporting soil protection policy.

At the core of the given definition of soil quality is the ability of soil to fulfil some important functions, and it is around this concept that the SQI literature has evolved. The SQI is a value, combining a variety of information regarding soil (chemical, physical and biological characteristics), which scores its 'fitness' to accomplish one or more functions (a low fitness results in a low SQI value). Interpretation of the overall 'soil health' or 'fitness for purpose' is thus simplified. Important soil functions include: water/solute flow and retention, physical stability and support, cycling of nutrients, filtration of potentially toxic materials, and maintenance of biodiversity and habitat (Daily et al., 1997). Typically, a SQI does not (and should not) cover all the above mentioned functions (Andrews et al., 2004): for a specific assessment, the functions of

* Corresponding author. Tel.: +39 080 5443004; fax: +39 080 5442976.
E-mail addresses: elena.armenise@gmail.com, e.armenise@cranfield.ac.uk (E. Armenise).

greatest relevance are selected according to the management goal for which the assessment is to be made.

Consequently, SQI assessment begins by defining the management goal(s) (e.g. crop productivity, waste recycling and environmental protection), followed by associating the critical soil functions. Since the functions are not directly measurable, appropriate physical, chemical and biological parameters, named soil quality indicators, are selected to indirectly measure how well each function is being performed (Karlen et al., 2004). The main requirement for a soil property to be selected as a soil quality indicator is that it shows sensitivity to changes occurring within the soil function in question. Other favourable features include: a positive correlation with ecosystem services, being easily measurable, sensitive to management, and whenever possible, to be a component of a pre-existing database (Andrews et al., 2004).

Various indexing techniques have been established so far (Andrews et al., 2002; Harris et al., 1996; Karlen and Stott, 1994; Wander and Bollero, 1999). The approach followed by Andrews et al. (2002), as based on statistical and mathematical tools, provides outcomes which are comparable across different soil types, regions and management systems. It was also the core upon which the Soil Management Assessment Framework (SMAF) was developed. SMAF is an additive, non-linear indexing tool, freely available for any user who desires to interpret soil quality indicators and integrate available information into an overall SQI (http://soilquality.org/tools/smaf_intro.html).

Soil management in an agricultural system can play a major role in soil protection. The success of any management in maintaining soil quality depends on how soil responds over time (Gregorich et al., 1994) and thus long-term field experiments provide an ideal platform for critical appraisal of these managements. Long-term experiments often produce huge data-sets, and handling such an amount of information can lead to confusion, and, in the worst instance, misinterpretation. SQIs can be used effectively for interpreting the multivariate data-set typical of long-term field trials (Hussain et al., 1999; Karlen et al., 1994, 2006; Masto et al., 2008; Sharma et al., 2008), with a single score being able to identify an overall trend among several, potentially conflicting indicators (Andrews et al., 2003).

In the present work, using data from an existing long-term field experiment comparing contrasting soil managements (cropping sequence, stubble management and NPK level), SQIs were calculated with the aim of quantifying the potential of the managements in upholding soil quality for agricultural purpose in the Mediterranean environment. The indexing technique assessed here follows that proposed by Andrews et al. (2002).

2. Materials and methods

2.1. Site description and soil analyses

This assessment of cropping sequence, stubble management and NPK-rate effects on SQI, used an existing long-term field study (38-year) located in Southern Italy (40°13'12" N, 16°40'48" E). The soil (FAO: Fluvisol, silty clay loam) general characteristics (0–0.35 m) in 2006 were 303 g clay kg⁻¹, 574 g silt kg⁻¹, 123 g sand kg⁻¹, 17.5 g organic C kg⁻¹, 60.3 g CaCO₃ kg⁻¹ and a pH 7.6 (soil:H₂O ratio 1:5).

The field trial is arranged in a split-split plot design with cropping sequence as the main plot, stubble management as the split plot, and fertiliser rate as the split-split plot. The treatments are replicated two times. Three wheat (*Triticum turgidum* L. var. *durum*) based crop sequences are compared: three-course rotation (sugar beet – wheat/dry bean double cropping – wheat [S-W₁/B-W₂]); continuous double cropping of wheat/dry bean [W/B]; continuous wheat [CW]. Eighteen NPK-rates and two stubble

managements (stubble incorporated [I] vs. burnt [B]) are also tested. However, for the purposes of this study, a reduced number of treatments were selected:

- wheat/dry bean double cropping in the 3-year rotation [W₁/B], continuous wheat in the 3-year rotation [W₂], continuous double cropping of wheat/dry bean [W/B], continuous wheat [CW];
- three NPK-rates: nil [F0], intermediate [F1], optimal [F2];
- stubble incorporated [I], burnt [B].

An adequate assessment of soil quality status through the SQI requires the contemporary analysis of as many soil parameters as possible in order to best represent the multifaceted nature of the soil. Parameters estimated in this study include 18 chemical and physical soil attributes: clay and sand content; gravimetric water content at –33, –800 and –1500 kPa matric potential (θ_{g33} , θ_{g800} and θ_{g1500}); plant available water (PAW); *S* index; particle density (ρ_r); soil organic matter (SOM); total N (TN); water extractable N and organic C (WEN, WEOC); soil pH; electrical conductivity (EC); available P (P_{av}); exchangeable Mg²⁺, K⁺, and Ca²⁺ (Mg_{ex}, K_{ex} and Ca_{ex}).

Composite samples were collected for soil tests from the selected field plots to a depth of 0–0.35 m using a 0.05 m diameter hand-auger. Water extractable organic C and water extractable N were determined on 2-mm sieved field-moist soil samples (soil water content = 0.107 g g⁻¹) collected in May 2007. Soil samples utilised for the remaining analyses were instead collected at the end of the wheat cropping cycle in July 2006, air-dried and crushed to pass a 2 mm sieve.

Soil pH and EC were determined at a 1:5 soil–water ratio with a glass electrode and conductivity bridge respectively. SOM was estimated by dichromate oxidation (Walkley and Black, 1934); TN by the Kjeldahl method (Bremner and Mulvaney, 1982). Available P was detected by sodium bicarbonate (NaHCO₃) extraction and subsequent colorimetric analysis (Olsen et al., 1954). Taking into account the soil pH (pH > 7), exchangeable Mg²⁺, K⁺, and Ca²⁺ were measured by displacement with barium chloride and quantification by atomic adsorption spectrometry (Hendershot and Duquette, 1986). Soil particle size distribution was measured using the pipette method after oxidation of the SOM with H₂O₂ and stirring in a sodium hexametaphosphate solution (Gee and Bauder, 1986). Soil particle density was determined by means of a pycnometer. WEOC and WEN were extracted from fresh soil according to a protocol obtained by combining procedures reported in Haynes (2000) and Rees and Parker (2005). Briefly, 30 g of fresh-weight soil and 60 ml distilled water (1:2, soil:H₂O) were placed in an Erlenmeyer flask, stoppered and shaken for 30 min. The extracts were centrifuged at 20,000 × g for 10 min and the supernatant was filtered through 0.45 μm Millipore filter. Total organic C and total N in the water extracts were analysed with a Shimadzu TOC/TN Analyzer (Model TOC-VCSH, Shimadzu, Kyoto, Japan).

The water-retention characteristics were investigated by means of pressure plate apparatus in the drying process (Klute, 1986). Disturbed soil samples, held in 2 cm high rings, were slowly wetted up to saturation using a sand table and then dried down to six distinct water potentials (–10, –33, –100, –500, –800 and –1500 kPa). After a period of equilibration, samples were removed and their water content determined gravimetrically. A smooth curve based on van Genuchten (1980) equation was fitted using RETC program (van Genuchten et al., 1991). This equation was chosen both because it provides a very good correlation between experimental and calculated data, and because its parameters could be used to compute a soil physical quality index, the *S* index (Dexter, 2004). *S* is the slope of soil water retention curve at its inflection point when the gravimetric water content (θ_g) is plotted

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات