



Differentiating hypochondriasis from panic disorder

Wolfgang Hiller^{a,b,*}, Rolf Leibbrand^a, Winfried Rief^c,
Manfred M. Fichter^{b,d}

^a*Psychological Institute, University of Mainz, Staudingerweg 9,
D-55099 Mainz, Germany*

^b*Roseneck Center of Behavioral Medicine, Prien, Germany*

^c*Psychological Institute, University of Marburg, Marburg, Germany*

^d*Medical Faculty, University of Munich, Munich, Germany*

Received 7 April 2003; received in revised form 22 October 2003; accepted 28 October 2003

Abstract

Hypochondriasis and panic disorder are both characterized by prevalent health anxieties and illness beliefs. Therefore, the question as to whether they represent distinct nosological entities has been raised. This study examines how clinical characteristics can be used to differentiate both disorders, taking the possibility of mixed symptomatologies (comorbidity) into account. We compared 46 patients with hypochondriasis, 45 with panic disorder, and 21 with comorbid hypochondriasis plus panic disorder. While panic patients had more comorbidity with agoraphobia, hypochondriasis was more closely associated with somatization. Patients with panic disorder were less pathological than hypochondriacal patients on all subscales of the Whiteley Index (WI) and the Illness Attitude Scales (IAS) except for illness behavior. These differences were independent of somatization. Patients with hypochondriasis plus panic had higher levels of anxiety, more somatization, more general psychopathology and a trend towards increased health care utilization. Clinicians were able to distinguish between patient groups based upon the tendency of hypochondriacal patients to demand unnecessary medical treatments. These results confirm that hypochondriasis and panic disorder are distinguishable

* Corresponding author.

E-mail address: hiller@mail.uni-mainz.de (W. Hiller).

clinical conditions, characterized by generally more psychopathology and distress in hypochondriasis.

© 2003 Elsevier Inc. All rights reserved.

Keywords: Hypochondriasis; Panic disorder; Health anxieties; Comorbidity; Somatization; Illness behavior

1. Introduction

Health anxieties are frequently experienced by patients with medical illness or mental disorders. They are perceived as plausible emotional reactions if a serious or life-threatening disease exists. In other cases, health anxieties may develop despite absence of organic pathology, especially when patients tend to misinterpret minor bodily sensations as signs of a serious disease or mistrust their doctors. If strong health anxieties persist over long periods of time and have negative consequences for the psychosocial functioning of the person, the diagnosis of hypochondriasis can be made. This diagnosis is included in categorical classification systems such as DSM-IV (American Psychiatric Association, 1994) and can be quantified by worldwide used hypochondriasis scales such as the Whiteley Index or the Illness Attitude Scales (Hiller, Rief, & Fichter, 2002).

However, hypochondriacal disorder is not the only clinical condition defined by predominant health anxieties. Patients with panic disorder usually report many intense somatic symptoms during their panic attacks such as palpitations and accelerated heart rate, shortness of breath, chest pain, nausea, paresthesia or dizziness. They also tend to attribute these symptoms to organic causes, such as heart or pulmonary disease. As a consequence, panic patients frequently demand extensive medical examinations or consult numerous specialists in the hope that the organic causes of their symptoms can be detected. Thus, the emotional, cognitive, and behavioral reactions of panic patients are very similar to those typically described for hypochondriacal disorder.

Despite these similarities, hypochondriacal and panic disorder can be well distinguished through use of structured interviews or diagnostic checklists (Barsky, Wyshak, & Klerman, 1992; Fava & Grandi, 1991). The major difference is the episodic nature of the symptoms in panic disorder versus the more or less persisting complaints in hypochondriasis. Panic patients usually experience their symptoms only during discrete periods that have a sudden onset and build to a peak within a few minutes, although worries concerning development of new attacks may persist in the intervals between the attacks. Hypochondriasis, on the other hand, is defined as fears or ideas of having a serious disease for more than 6 months. The exclusion criterion for hypochondriasis in DSM-IV (F) specifies that the disorder is not to be diagnosed if the symptomatology is fully accounted for by panic disorder. However, this does not generally exclude co-existence of hypochondriasis and panic disorder because patients may suffer from both episodic panic attacks

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات