



Changing the perspective on early development of Rett syndrome

Peter B. Marschik^{a,b,1}, Walter E. Kaufmann^{c,1,*}, Jeff Sigafos^d, Thomas Wolin^b, Dajie Zhang^e,
 Katrin D. Bartl-Pokorny^b, Giorgio Pini^f, Michele Zappella^f, Helen Tager-Flusberg^g,
 Christa Einspieler^b, Michael V. Johnston^h

^a Center for Genetic Disorders of Cognition and Behavior, Kennedy Krieger Institute, Department of Neurology, Johns Hopkins University School of Medicine, Baltimore, USA

^b Institute of Physiology, Center for Physiological Medicine, Medical University of Graz, Graz, Austria

^c Department of Neurology, Boston Children's Hospital and Harvard Medical School, Boston, USA

^d Department of Educational Psychology, Victoria University of Wellington, Wellington, New Zealand

^e Department of Psychology, Karl-Franzens University of Graz, Graz, Austria

^f Tuscany Rett Centre Versilia Hospital, Lido di Camaiore, Italy

^g Department of Psychology, Boston University, Boston, USA

^h Department of Neurology and Pediatrics, Johns Hopkins University School of Medicine, Kennedy Krieger Institute, Baltimore, USA

ARTICLE INFO

Article history:

Received 18 January 2013

Accepted 18 January 2013

Available online 9 February 2013

Keywords:

Rett syndrome

Preserved speech variant

Speech-language development

Milestones

Video analysis

Regression

ABSTRACT

We delineated the achievement of early speech-language milestones in 15 young children with Rett syndrome (*MECP2* positive) in the first two years of life using retrospective video analysis. By contrast to the commonly accepted concept that these children are normal in the pre-regression period, we found markedly atypical development of speech-language capacities, suggesting a paradigm shift in the pathogenesis of Rett syndrome and a possible approach to its early detection.

© 2013 Elsevier Ltd. Open access under [CC BY-NC-ND license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

1. Introduction

Mutations in the X-linked gene encoding Methyl-CpG-binding protein 2 (MeCP2) account for 95–97% of cases of Rett syndrome (RTT, MIM 312750), a neurodevelopmental disorder affecting predominantly females (Amir et al., 1999; Hagberg, Aicardi, Dias, & Ramos, 1983; Neul et al., 2010). Key features of typical RTT, and the relatively milder preserved speech variant (PSV), are stereotyped hand movements coinciding with a regression in purposeful hand use and expressive language (Hagberg et al., 1983; Kerr, Archer, Evans, & Gibbon, 2006; Matson, Fodstad, & Boisjoli, 2008; Neul et al., 2010; Renieri et al., 2009). The dynamic course of RTT is said to involve a period of apparently normal early development followed by a profound neurologic regression and subsequent stabilization or partial recovery. However, this view contrasts with growing

* Corresponding author at: Department of Neurology, Boston Children's Hospital and Harvard Medical School, 300 Longwood Avenue, LO 462, Boston, MA 02115, USA. Tel.: +1 617 355 0537.

E-mail address: walter.kaufmann@childrens.harvard.edu (W.E. Kaufmann).

¹ These authors contributed equally to this work.

knowledge about *MECP2* expression in the developing brain and its role in terminal neuronal differentiation (Kaufmann, Johnston, & Blue, 2005; Zoghbi, 2003). Over the last two decades there has been a shift from some initial thoughts about atypical pre-regressional development in individuals with RTT to the growing evidence of early behavioral abnormalities reflecting a poor integration of the young nervous system (Einspieler, Kerr, & Prechtl, 2005a, 2005b; Leonard & Bower, 1998; Marschik, Einspieler, Oberle, Laccone, & Prechtl, 2009; Marschik, Einspieler, Prechtl, Oberle, & Laccone, 2010; Marschik, Einspieler, & Sigafos, 2012; Marschik, Kaufmann, et al., 2012; Tams-Little & Holdgrafer, 1996). The redefined clinical diagnostic criteria by the RettSearch consortium (Neul et al., 2010) highlight the age-dependency of clinical changes and suggest the existence of initial signs in RTT, which might appear earlier than, for example, the (non obligatory) deceleration of head growth. Nonetheless, the nature of these early signs and their potential for early and reliable identification of RTT are still unclear. Consequently, the present study aimed at shedding light into pre-regressional development in RTT by focusing on early speech-language development using a video database.

2. Materials and methods

Audio-video recordings of sufficient length and quality standards during the first two years of life were available for 15 RTT participants with *MECP2* mutations (10 typical RTT, 5 PSV; Neul et al., 2010; Renieri et al., 2009) comprising a total footage of 2431 min (i.e., 40.5 h; typical RTT 2041 min, median = 68; PSV 390 min, median = 78). All RTT participants were females, born as a singleton after uneventful pregnancy and delivery; birth weights, birth lengths, occipitofrontal circumferences, and Apgar scores were within the normal ranges. The recordings of play situations, daily routines and special events were made by the parents at a time when they were not aware that their daughters had typical RTT or PSV. The recordings were copied, prepared for analyses by unifying codecs, sampled across the age range and analyzed retrospectively.

We focused on and coded the achievement and realization of age-specific speech-language milestones. These included: protophones such as cooing and babbling vocalizations, proto-words, word combinations, and the communicative milestone of intentional gestures (Fig. 1). In addition to the achievement of these milestones, we also focused on the detection of atypical vocalizations during the cooing and babbling period. Fixed vocal signals (e.g., crying, laughing), vegetative sounds (e.g., sneezing, coughing), and utterances unintelligible to the coders were excluded. All remaining vocalizations were transcribed in chronological order and analyses were carried out using the Noldus Observer-XT (Versions 9.0 and 11.0, Noldus Information Technology, <http://www.noldus.com>). Two PhD students (TW, KDBP; blind to the background and purpose of the study) and two of the authors (PBM, CE) coded the data in parallel with an inter-rater agreement of 97%. The study was approved by the relevant Institutional Review Boards and all parents gave their informed consent, including consenting to the publication of the results.

3. Results

We focused on the extent to which children reached developmental milestones in the speech-language domain and found that, with each sequential milestone, more participants failed to reach the next level of complexity. Whereas 13/15 participants, including all of those with PSV, showed cooing vocalizations only 9 had canonical babbling (half of the females with typical RTT and all but one with PSV; Fig. 1). Proto-words were largely absent in participants with typical RTT (only 3/10), although they were observed in three participants with PSV; furthermore, none of the participants with typical RTT and only one of those with PSV produced any word combinations (Fig. 1). Only 5/10 typical RTT and 3/5 PSV participants actively used communicative gestures.

Qualitative analyses of verbal behaviors revealed frequent abnormalities, even when the milestones of cooing and babbling were reached. Vocalizations on inspiratory airstream were among the first atypical features observed during the cooing and babbling period in 7/10 typical RTT and 2/5 PSV participants. Normal cooing vocalizations were observed in 5/10 typical RTT participants whereas 2/10 exhibited an intermittent character of normal and abnormal cooing (i.e., typical age-specific vocalizations interspersed with atypical ones such as inspiratory vocalizations, pressed or high-pitched crying-like vocalizations), and 3/10 did not display any cooing vocalizations. For PSV, 3/5 participants had normal and 2/5 had




Fig. 1. Step by step: Trajectories of reaching developmental speech-language milestones in typical RTT ($n = 10$) or PSV ($n = 5$) during the first two years of life.

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات