



unequivocally demonstrated that the MTL is indispensable for conscious memory in humans (Eichenbaum, 2013; Morris, 2007; Squire and Wixted, 2011; Stark, 2007). However, there is a continuing debate on the precise contributions of the MTL and its subregions to distinct memory domains and to cognition beyond memory such as perception, decision making and imagination of the future (Henke, 2010; Ranganath, 2010; Shohamy and Turk-Browne, 2013; Squire and Wixted, 2011). The current lack of a unifying framework for the role of the MTL in memory and other cognitive functions may also be due to the fact that human lesion models of MTL dysfunction differ considerably with respect to the temporal properties, selectivity, extent and laterality of lesions (Stark, 2007). In addition, only very few patients with autopsically verified selective bilateral lesions to the MTL have been reported (Zola-Morgan et al., 1986; Rempel-Clower et al., 1996). In many cases, significant uncertainties remain about the functional status of brain regions in- and outside the MTL (Squire and Wixted, 2011; Stark, 2007). Beside differences in assessment of memory between studies, these factors may significantly influence performance in cognitive tasks. For example, we have shown recently that the temporal properties of disorders affecting the MTL critically determine a subjects' performance in visuo-spatial short-term memory tasks, even in patients with otherwise similar lesion characteristics (Braun et al., 2008; Finke et al., 2013). Although previous studies in other domains such as motor function and language have repeatedly demonstrated a significant influence of disease type on behavioral performance (e.g. Anderson et al., 1990; Haaland and Delaney, 1981), systematic investigations on how mechanisms of MTL damage may account for the partly divergent findings in human lesion studies of memory have only rarely been reported.

Here, we have investigated the role of disease characteristics for behavioral performance in memory tasks. Patients with MTL lesions or MTL dysfunction acquired in the context of three different disorders (benign brain tumor, herpes encephalitis, global cerebral hypoxia) were tested with a set of tasks that included testing for short-term memory of colors, locations and color–location associations. Consistent with the hypothesis that the hippocampus is particularly involved in associative binding (Henke, 2010; Ranganath, 2010; Aggleton et al., 2012; Yonelinas, 2013), patients with damage to the hippocampal formation have previously shown selective deficits in memory of color–location associations, while performance in the other tasks was normal (Braun et al., 2008, 2011; Finke et al., 2008). In the present study, deficient associative memory in all patients suggests that all patient groups may provide appropriate models of hippocampal dysfunction. However, the presence of presumably MTL-independent deficits in non-associative memory tasks appears to depend significantly on disease characteristics.

## 2. Methods

### 2.1. Subjects

Thirty-one patients were recruited from the Department of Neurology and the Department of Nephrology and Medical Intensive Care Medicine at the Charité-Universitätsmedizin Berlin, Germany (6 patients with right-sided MTL lesions following resection of a benign brain tumor, 5 patients with left-sided MTL lesions following resection of a benign brain tumor, 10 patients with MTL lesions following herpes simplex encephalitis and 10

**Table 1**  
Patient characteristics.

Patient	Diagnosis	Sex	Age	Delay (months) since lesion	Clinical notes/Histopathology	Anticonvulsant medication/centrally acting drugs
1	R-MTL	F	42	22	Pilomyxoid astrocytoma	Gabapentin 900 mg/d
2	R-MTL	F	32	44	Epidermoid tumor	Lamotrigine 200 mg/d
3	R-MTL	M	19	47	Pilocytic astrocytoma	None
4	R-MTL	M	24	5	Neuroepithelial tumor	Lamotrigine 200 mg/d
5	R-MTL	M	24	56	Pigmented astrocytoma	None
6	R-MTL	F	23	21	Ganglioglioma	Levetiracetam 2 g/d
7	L-MTL	M	22	8	DNET tumor WHO I	Oxcarbazepine 1800 mg/d
8	L-MTL	F	35	113	Ganglioneuroma	None
9	L-MTL	F	45	84	Oligo-astrocytoma WHO II	None
10	L-MTL	F	49	2	Ganglioglioma	Levetiracetam 3 g/d; lamotrigine 25 mg/d
11	L-MTL	M	38	18	Cavernous Haemangioma	Levetiracetam 2 g/d
12	HSE	M	33	100	Herpes 1 PCR positive	Phenytoin 500 mg/d
13	HSE	F	61	24	Herpes 1 PCR positive	Carbamazepine 600 mg ret./d
14	HSE	M	35	96	Herpes 1 PCR positive	None
15	HSE	F	69	28	Herpes 1 PCR positive	None
16	HSE	F	54	96	Herpes 1 PCR positive	Levetiracetam 3 g/d; oxcarbazepine 1200 mg/d; escitalopram 20 mg/d
17	HSE	M	32	6	Herpes PCR neg./HSV IgM in CSF positive	Doxepin 100 mg/d
18	HSE	M	68	3	Herpes 1 PCR positive	Levetiracetam 2 g/d
19	HSE	M	49	5	Herpes 1 PCR positive	Levetiracetam 2 g/d
20	HSE	F	67	14	Herpes 1 PCR positive	None
21	HSE	M	53	2	Herpes 1 PCR positive	Citalopram 20 mg/d; opipramol 150 mg/d
22	GCH	M	27	95	CPC score 2, no hypothermia	None
23	GCH	M	69	36	CPC score 1, therapeutic hypothermia	None
24	GCH	M	57	30	CPC score 1, therapeutic hypothermia	None
25	GCH	M	57	33	CPC score 2, therapeutic hypothermia	None
26	GCH	M	35	36	CPC score 1, therapeutic hypothermia	None
27	GCH	M	54	23	CPC score 1, therapeutic hypothermia	None
28	GCH	M	68	11	CPC score 1, no hypothermia	None
29	GCH	F	62	10	CPC score 2, therapeutic hypothermia	None
30	GCH	M	60	8	CPC score 2, therapeutic hypothermia	None
31	GCH	F	52	5	CPC score 1, therapeutic hypothermia	Lorazepam 0,5 mg/d; citalopram 20 mg/d

متن کامل مقاله

دریافت فوری ←

**ISI**Articles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات