

Accepted Manuscript

Improvements of ambient hyperglycemia and glycemic variability are associated with reduction in oxidative stress for patients with type 2 diabetes

Makoto Ohara, Hiroe Nagaïke, Satoshi Goto, Ayako Fukase, Yuki Tanabe, Masako Tomoyasu, Takeshi Yamamoto, Toshiyuki Hayashi, Tomoyasu Fukui, Tsutomu Hirano


PII: S0168-8227(17)31645-5
DOI: <https://doi.org/10.1016/j.diabres.2018.02.017>
Reference: DIAB 7230

To appear in: *Diabetes Research and Clinical Practice*

Received Date: 1 November 2017
Revised Date: 31 January 2018
Accepted Date: 13 February 2018

Please cite this article as: M. Ohara, H. Nagaïke, S. Goto, A. Fukase, Y. Tanabe, M. Tomoyasu, T. Yamamoto, T. Hayashi, T. Fukui, T. Hirano, Improvements of ambient hyperglycemia and glycemic variability are associated with reduction in oxidative stress for patients with type 2 diabetes, *Diabetes Research and Clinical Practice* (2018), doi: <https://doi.org/10.1016/j.diabres.2018.02.017>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Improvements of ambient hyperglycemia and glycemic variability are associated with reduction in oxidative stress for patients with type 2 diabetes

Makoto Ohara^a, Hiroe Nagaike^a, Satoshi Goto^a, Ayako Fukase^a, Yuki Tanabe^a, Masako Tomoyasu^a, Takeshi Yamamoto^a, Toshiyuki Hayashi^a, Tomoyasu Fukui^a, Tsutomu Hirano^a

^a Department of Medicine, Division of Diabetes, Metabolism, and Endocrinology, Showa University School of Medicine, Tokyo, Japan

Correspondence to Makoto Ohara, MD, PhD

1-5-8 Hatanodai, Shinagawa-ku, Tokyo 142-8666, Japan

Tel: +81-3-3784-8000

Fax: +81-3-3784-8948

E-mail: s6018@nms.ac.jp

Highlights

- Improvements in daily and day-to-day glucose and oxidative stress are associated.
- Glucose-like peptide 1 receptor agonists directly affect oxidative stress.
- Statin relieves oxidative stress independent of improvements in lipids.

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات