


Accepted Manuscript

Solubility of taurine and its application for the crystallization process improvement

Di Wu, Liangcheng Song, Chongqiang Zhu, Xiao Zhang, Huai Guo, Chunhui Yang



PII: S0167-7322(17)31744-0
DOI: doi: [10.1016/j.molliq.2017.06.043](https://doi.org/10.1016/j.molliq.2017.06.043)
Reference: MOLLIQ 7489

To appear in: *Journal of Molecular Liquids*

Received date: 23 April 2017

Revised date: 4 June 2017

Accepted date: 10 June 2017

Please cite this article as: Di Wu, Liangcheng Song, Chongqiang Zhu, Xiao Zhang, Huai Guo, Chunhui Yang, Solubility of taurine and its application for the crystallization process improvement, *Journal of Molecular Liquids* (2017), doi: [10.1016/j.molliq.2017.06.043](https://doi.org/10.1016/j.molliq.2017.06.043)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Solubility of Taurine and Its Application for the Crystallization Process Improvement

Di Wu, Liangcheng Song^{*}, Chongqiang Zhu, Xiao Zhang, Huai Guo, Chunhui Yang

School of Chemistry and Chemical Engineering, Harbin Institute of Technology, Harbin 150001, People's Republic of China

ABSTRACT

The solubility of taurine in the binary water + acetone solvent mixture at temperatures from 278.15K to 323.15K was measured using the laser monitoring technique, and the experimental data were well correlated by the Jouyban–Acree model. The solubility decreased quickly when the mole fraction of acetone increased in the binary solvent mixture and also the metastable zone width got narrowed at the same time. Then the nucleation mechanism was investigated based on the induction time. The results showed that the homogeneous dominated at higher supersaturation while the heterogeneous nucleation dominated at lower supersaturation. Finally, a combined cooling and anti-solvent crystallization process was put forward. The product yield was raised to 96% from 73%, compared with the original cooling crystallization process in aqueous solution. What's more, the caking of taurine crystals was reduced and the fluidity of the product obtained was improved.

KEYWORDS:

Taurine; Solubility; Combined cooling and anti-solvent crystallization process; Nucleation mechanism; Metastable zone width.

^{*} To whom correspondence should be addressed.

E-mail: lcsong@hit.edu.cn; Phone: +86-451-86403829; Fax: +86-451-86418270.

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات