
### Accepted Manuscript

Weak solutions to non-homogeneous boundary value problems for time-fractional diffusion equations

Masahiro Yamamoto



 PII:
 S0022-247X(17)31049-1

 DOI:
 https://doi.org/10.1016/j.jmaa.2017.11.048

 Reference:
 YJMAA 21847

To appear in: Journal of Mathematical Analysis and Applications

Received date: 16 April 2016

Please cite this article in press as: M. Yamamoto, Weak solutions to non-homogeneous boundary value problems for time-fractional diffusion equations, *J. Math. Anal. Appl.* (2018), https://doi.org/10.1016/j.jmaa.2017.11.048

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Weak solutions to non-homogeneous boundary value problems for time-fractional diffusion equations

Masahiro Yamamoto \*

#### Abstract

We discuss an initial-boundary problem for a time-fractional diffusion equation with non-zero Dirichlet boundary values which belong to  $L^2$  in time t and to a Sobolev space of negative order in space and prove the unique existence of weak solutions and a priori estimates. The proof is based on the Caputo fractional derivative in Sobolev spaces and the transposition method. We show one application to the existence of solution to an optimal control problem.

#### §1. Introduction

Let  $\Omega \subset \mathbb{R}^n$  be a bounded domain with boundary  $\partial \Omega$  of  $C^2$ -class and let  $\nu = \nu(x) = (\nu_1, ..., \nu_n)$  be the unit outward normal vector to  $\partial \Omega$  at x. In this paper we consider an initial-boundary problem for a time-fractional diffusion:

$$\partial_t^{\alpha} u(x,t) = -Lu(x,t), \quad x \in \Omega, \ 0 < t \le T,$$
(1.1)

$$u(x,t) = g(x,t), \quad x \in \partial\Omega, \ 0 < t < T,$$
(1.2)

$$u(x,0) = 0, \quad x \in \Omega. \tag{1.3}$$

Throughout this paper, we assume  $0 < \alpha < 1$ , and by  $\partial_t^{\alpha} u$ , we denote the Caputo derivative, which can be defined for  $w \in C^1[0,T]$  by

$$\partial_t^{\alpha} w(t) = \frac{1}{\Gamma(1-\alpha)} \int_0^t (t-s)^{-\alpha} \frac{dw}{ds}(s) ds, \quad 0 < t \le T, \quad 0 < \alpha < 1, \tag{1.4}$$

\*Department of Mathematical Sciences, The University of Tokyo, Komaba, Meguro, Tokyo 153, Japan, e-mail: myama@ms.u-tokyo.ac.jp

# دريافت فورى 🛶 متن كامل مقاله

- امکان دانلود نسخه تمام متن مقالات انگلیسی
   امکان دانلود نسخه ترجمه شده مقالات
   پذیرش سفارش ترجمه تخصصی
   امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
   امکان دانلود رایگان ۲ صفحه اول هر مقاله
   امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
   دانلود فوری مقاله پس از پرداخت آنلاین
   پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات
- ISIArticles مرجع مقالات تخصصی ایران