

Accepted Manuscript

Combination of Separation Methods and Data Mining Techniques for Prediction of Anomalous Areas in Susanvar, Central Iran

Seyyed Saeed Ghannadpour, Ardeshir Hezarkhani, Taraneh Roodpeyma



PII: S1464-343X(17)30301-1

DOI: [10.1016/j.jafrearsci.2017.07.015](https://doi.org/10.1016/j.jafrearsci.2017.07.015)

Reference: AES 2967

To appear in: *Journal of African Earth Sciences*

Please cite this article as: Seyyed Saeed Ghannadpour, Ardeshir Hezarkhani, Taraneh Roodpeyma, Combination of Separation Methods and Data Mining Techniques for Prediction of Anomalous Areas in Susanvar, Central Iran, *Journal of African Earth Sciences* (2017), doi: 10.1016/j.jafrearsci.2017.07.015

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Combination of Separation Methods and Data Mining Techniques for Prediction of Anomalous Areas in Susanvar, Central Iran

Seyyed Saeed Ghannadpour¹, Ardeshir Hezarkhani^{1,*}, Taraneh Roodpeyma¹

1- Department of Mining and Metallurgical Engineering, Amirkabir University of Technology (Tehran Polytechnic), 424 Hafez Ave., Tehran 15875-4413, Iran

Abstract

Structural method U-statistics is an eminent technique for delineating geochemical patterns; on the other hand, it is worthwhile to introduce Mahalanobis distance approach decreasing the background effects and intensifying the correlation factor of points as a powerful non-structural method. Undoubtedly, predicting the anomalous values could play an important role in the inchoate stages of exploration. Therefore, it is essential to find the most accurate approach to separate anomalous values from background and afterward use the results to anticipate each arbitrary sample. In this study, results of the combination between U-statistics & Mahalanobis distance algorithms are used to distinguish anomalous values from background on an accurate point of view. Then, three data mining methods will be applied to produce practical equations and finally determine anomalous values. Separation of geochemical anomalies, based on the combination of the U-statistics and the Mahalanobis distance approaches, would be done; then, under the influence of their results and the other parameters – x and y coordinates and Au and As grades - three data mining methods, K nearest neighbor (K-NN), decision tree, and naïve Bayes classifier, have been applied. For this purpose after separation of anomalous values according to the number of 603 samples by applying above combination, the data mining methods would be utilized to anticipate

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات