

# Accepted Manuscript

Self-quenched semiconducting polymer nanoparticles for amplified *in vivo* photoacoustic imaging

Chen Xie, Paul Kumar Upputuri, Xu Zhen, Manojit Pramanik, Kanyi Pu



PII: S0142-9612(16)30693-7

DOI: [10.1016/j.biomaterials.2016.12.004](https://doi.org/10.1016/j.biomaterials.2016.12.004)

Reference: JBMT 17853

To appear in: *Biomaterials*

Received Date: 14 August 2016

Revised Date: 14 October 2016

Accepted Date: 6 December 2016

Please cite this article as: Xie C, Upputuri PK, Zhen X, Pramanik M, Pu K, Self-quenched semiconducting polymer nanoparticles for amplified *in vivo* photoacoustic imaging, *Biomaterials* (2017), doi: 10.1016/j.biomaterials.2016.12.004.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

# Self-Quenched Semiconducting Polymer Nanoparticles for Amplified In Vivo Photoacoustic Imaging

Chen Xie, Paul Kumar Upputuri, Xu Zhen, Manojit Pramanik and Kanyi Pu\*

School of Chemical and Biomedical Engineering, Nanyang Technological University,  
Singapore 637457, Singapore

Address correspondence to: kypu@ntu.edu.sg

## Abstract

Development of photoacoustic (PA) imaging agents provides opportunities for advancing PA imaging in fundamental biology and medicine. Despite the promise of semiconducting polymer nanoparticles (SPNs) for PA imaging, the molecular guidelines to enhance their imaging performance are limited. In this study, semiconducting polymers (SPs) with self-quenched fluorescence are synthesized and transformed into SPNs for amplified PA imaging in living mice. The self-quenched process is induced by the incorporation of an electron-deficient structure unit into the backbone of SPs, which in turn promotes the nonradiative decay and enhances the heat generation. Such a simple chemical alteration of SP eventually leads to 1.7-fold PA amplification for the corresponding SPN. By virtue of the targeting capability of cyclic-RGD, the amplified SPN can effectively delineate tumor in living mice and increase the PA intensity of tumor by 4.7-fold after systemic administration. Our study thus provides an effective molecular guideline to amplify the PA brightness of organic imaging agents for *in vivo* PA imaging.

**Keywords:** Photoacoustic imaging; Semiconducting polymer nanoparticles; Tumor imaging; Contrast agents

متن کامل مقاله

دریافت فوری ←

**ISI**Articles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات