



The effect of pleasant olfactory mental imagery on the incidence and extent of atelectasis in patients after open heart surgery

Masoud Rezaei-Nodehi^a, Seyed Afshin Shorofi^{b,c}, Masoumeh Bagheri-Nesami^{d,*},
Rahman Ghafari^e, Jamshid Yazdani-Charati^f, Manizheh Darbeheshti^g

^a Student Research Committee, School of Nursing and Midwifery, Addiction Institute, Mazandaran University of Medical Sciences, Sari, Iran

^b Traditional and Complementary Medicine Research Center, Addiction Institute, Mazandaran University of Medical Sciences, Sari, Iran

^c Adjunct Research Fellow, Flinders University, Adelaide, Australia

^d Traditional and Complementary Medicine Research Center, Addiction Institute, Mazandaran University of Medical Sciences, Sari, Iran

^e Department of Cardiac Surgery, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

^f Health Sciences Research Center, Mazandaran University of Medical Science, Sari, Iran

^g Department of Radiology, Mazandaran Cardiac Surgery Center, Sari, Iran

ARTICLE INFO

Keywords:

Mental imagery
Olfactory imagery
Atelectasis
Heart surgery

ABSTRACT

Background and purpose: Atelectasis is the most common pulmonary complication after open heart surgery. This study was intended to examine the effects of pleasant olfactory mental imagery on postoperative atelectasis in patients undergoing open heart surgery.

Materials and methods: This is a randomized controlled clinical trial. The sample consisted of 80 patients who were randomly assigned to either practice olfactory mental imagery (test group) or receive routine care (control group). A card with the image of roses was given to patients and they were asked to look at the image, visualize the scent of roses in the mind, and then sniff as much as possible, hold their breath for 2 s and eventually exhale slowly through the nose. This procedure was consecutively repeated five times. After a fifteen-minute break, patients proceeded to practice olfactory mental imagery with other fruit images (banana, apple, and lemon). The test group executed the olfactory mental imagery for two hours in the morning and two hours in the afternoon on postoperative days 1 and 2. The control group received the routine ICU care. A questionnaire collected information on sociodemographic characteristics and clinical parameters. Chest radiographs were used to diagnose atelectasis, which were evaluated by the hospital radiologist.

Results: No statistically significant differences were observed between the two groups regarding socio-demographic, medical and surgical information. The incidence of atelectasis in the test group (40%, n = 16) was significantly lower than in the control group (67.5%, n = 27) on postoperative day 2 (p = 0.02).

Conclusion: Our findings suggest that olfactory mental imagery can improve respiratory function and reduce the risk of atelectasis in patients with cardiac surgery.

1. Introduction

Early pulmonary complications are common after cardiac surgery,^{1–3} such complications include atelectasis, pneumonia, pleural effusion, and acute respiratory distress syndrome.^{4,5} Atelectasis is the most common pulmonary complication after open heart surgery⁵ with the incidence of 15–98% after heart and thoracic surgeries.⁶ In Iran, the incidence of atelectasis has been reported to be as high as 75% and 56.7% following thoracic⁷ and coronary artery bypass surgeries,⁸ respectively. Atelectasis is the collapse of a group of alveoli, a small lobe or larger lung units, which can be caused by a variety of factors

following surgery (specifically thoracic surgery) including external cardiopulmonary bypass, sedative medications,^{9,10} anesthesia,^{3,10,11} respiratory muscle dysfunction,^{11–13} pain from the sternotomy, pain from the chest drains,^{10,14–16} impaired phrenic nerve function, and diaphragmatic dysfunction.^{5,11,17,18} There may be no significant clinical symptoms of post-surgical atelectasis. However, post-surgical atelectasis has a progressive process and may contribute to serious pulmonary complications including nosocomial pneumonia, which can increase the length of hospital stay and health care costs.^{1,19–21}

Despite the use of active (deep breathing along with coughing, incentive spirometry, frequent repositioning in the bed, and early

* Corresponding author.

E-mail address: anna30432003@yahoo.com (M. Bagheri-Nesami).

ambulation) and passive (intermittent positive-pressure breathing, positive end expiratory pressure, non-invasive positive-pressure ventilation) respiratory physiotherapy techniques following cardiac surgery,²² the incidence of post-surgical atelectasis is still considerable.^{7,23} There is also insufficient evidence to support the effectiveness of respiratory physiotherapy techniques on the prevention of pulmonary complications after thoracic surgery.^{6,24–28}

Olfactory mental imagery is the ability to sense a smell in the absence of any olfactory stimulus and is actually a short memory of olfactory events with the mind's nose.²⁹ Olfactory mental imagery is thought to involve the same brain areas as olfactory perception.^{30–32} Similar to odor perception allowed by the sense of smell, olfactory mental imagery gives also rise to sniffing,³¹ which does not depend on the power of individuals' imagination.^{33,34} Several studies indicated that pleasant odors are sniffed stronger and longer than unpleasant odors.^{34–36} One of the characteristics of the olfactory system is its close connection with the respiratory system.^{36,37} Kleeman et al. found that “changes in the breathing profile result from sniffing that accompanied both olfactory perception and olfactory imagery” (p.8). They also explored that respiratory minute volume and respiratory amplitude increase during pleasant olfactory imagery in comparison to baseline. This increase in respiratory parameters is brought about by the elevated tidal volume and the improved breathing pattern and is not associated with the respiratory rate.³³ There are two assumptions about the improvement of respiratory function following olfactory mental imagery. The first one is that breathing pattern is often slow and deep with the least rate during the feelings of comfortableness and pleasantness. In other words, deep breathing and pleasantness mutually affect each other. Pleasant olfactory mental imagery contributes to increased feelings of pleasantness and comfortableness, leading to deep breathing and reduced respiratory rate.³⁸ The second hypothesis suggests that there is a deep synchronization between the brain's slow-wave (alpha waves) across the entire cortical mantle and slow and deep breathing pattern during sleep. “Humans have learned, through the conscious control of slow breathing, to induce a level of whole-brain synchronization that is naturally found only during slow-wave sleep. However, when induced in the active state, the synchronization occurs at the higher frequencies associated with controlled mentation” (p.435).³⁹ Pleasant olfactory mental imagery can also stimulate lung baroreceptors through deep breathing caused by the sniff reflex. Stimulating these receptors leads to an increase in the frequency and duration of inhibitory neural impulses, resulting in parasympathetic dominance. Alpha waves are also synchronized through enhanced activation of the parasympathetic system.⁴⁰ Thus, deep breathing and alpha waves mutually affect each other, which facilitates deep breathing pattern.

It has been shown that odor-evoked autobiographical pleasant memory retrieval may extend the inspiratory and expiratory time (particularly the expiratory time) through pleasantness of the memory or increased parasympathetic tone, resulting in deep breathing and reduced respiratory rate.³⁸ This respiratory pattern is not only controlled by voluntary deep breathing and changes in metabolic demand, but also it is affected by olfactory-related brain regions.⁴¹ Prolonged expiration triggered by olfactory imagery³⁸ increases intra-thoracic pressure,⁴² which pushes more blood into the heart with a subsequent increase in stroke volume. Baroreceptors in aortic arch receive more pressure caused by the increased stroke volume, which inhibits the tonic activity in vasoconstrictor fibers and excites the vagal innervations of the heart with resultant vasodilatation, a decrease in blood pressure and bradycardia.^{40,42} Vagal cardiac and pulmonary mechanisms are linked, thereby the changes in the cardiovascular system may have effects on the respiratory system.³⁸ Therefore, an increase in pulmonary vagal tone leads to slow and deep breathing pattern.

Odor adaptation does not take place after olfactory imagery as the physical odor stimulus does not exist. Therefore, olfactory imagery can be used repeatedly in session. On the other hand, given that the cerebellum regulates inversely the sniff volume in relation to odor

concentration,^{33,43} maximum sniffs occur during olfactory imagery with physically absent odors. Considering that nasal deep breathing more than oral deep breathing can improve respiratory function,⁴⁴ it seems that pleasant olfactory mental imagery and resultant deep inspirations in the form of sniffing can become an easy part of respiratory exercise.

To the best of our knowledge, no published study has explored the effect of pleasant olfactory mental imagery on postoperative atelectasis in patients undergoing open heart surgery. Given the high incidence of atelectasis following open heart surgery, despite active and passive breathing exercises, and the role of olfactory mental imagery in inducing deep breathing and the expansion of lung volumes, this study was intended to examine the effects of pleasant olfactory mental imagery on postoperative atelectasis in patients undergoing open heart surgery.

2. Materials and methods

This is a randomized controlled clinical trial. The study population included all patients undergoing open heart surgery at a hospital affiliated to the Mazandaran University of Medical Sciences, Sari, Iran. The sample consisted of 80 patients who were consecutively enrolled and randomly assigned to either practice olfactory mental imagery (test group) or receive routine care (control group) according to a computer-generated randomization list. The eligibility criteria included patients scheduled for elective, non-emergency, open heart surgery; age 18 years and older^{45,46}; lack of mental impairment; ejection fraction higher than 30 percent measured with echocardiography prior to cardiac surgery owing to its significant impact on the respiratory system;⁴⁷ no previous heart or lung surgery; lack of any chronic pulmonary diseases such as atelectasis and pneumonia according to treating doctors¹⁷; no previous severe head or nasal injuries; and no previous neurological disorders and recurrent sinus infections.³⁴ In addition, patients were not enrolled in the study if they had systolic blood pressure lower than 90 mmHg despite adequate fluid intake, arterial blood pH less than 7.30, partial pressure of arterial CO₂ higher than 50 mmHg, arterial oxygen saturation less than 80% even with breathing supplemental oxygen, hemoglobin level less than 7 g/dL, serum creatinine level higher than 3.5 mg/dL, and a body mass index (BMI) of more than 40 kg/m² due to the increased risk of respiratory complications after surgery.⁴⁸ The exclusion criteria were a decision to withdraw from the study, postoperative hemodynamic instability (systolic blood pressure less than 80 mmHg),⁷ postoperative bleeding more than 500 ml in the first postoperative hour,²⁰ and the need for mechanical ventilatory support for more than 15 h after surgical operation.²

All patients signed an informed consent form if agreed to participate in the study. To accustom participants to the experiment, patients in the test group underwent training for the olfactory mental imagery on the day before surgery. A card with the image of roses was given to patients and they were asked to look at the image, visualize the scent of roses in the mind, and then sniff as much as possible, hold their breath for 2 s and eventually exhale slowly through the nose. This procedure was repeated five times for image of roses. After a fifteen-minute break, patients proceeded to practice olfactory mental imagery with other fruit images (banana, apple, and lemon) as similar as with the image of roses. In order to facilitate secretion clearance from the airways, patients were to be encouraged to have two effective coughs during exhalation following five consecutive olfactory mental imageries for each image. Each patient was expected to perform a total of twenty imagery tasks in an hour.

After the training session, the test group executed the olfactory mental imagery for two hours in the morning and two hours in the afternoon on postoperative days 1 and 2. The control group received the routine ICU care. A questionnaire collected information on socio-demographic characteristics and clinical parameters including age, sex, and marital status, highest level of educational attainment, smoking status and BMI. The questionnaire also recorded information on type of

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات