

## Author's Accepted Manuscript

Early maladaptive schemas and suicidal risk in an Iranian sample of patients with obsessive-compulsive disorder

Vahid Khosravani, Farangis Sharifi Bastan, Mehdi Samimi Ardestani, Razieh Jamaati Ardakani



PII: S0165-1781(16)31811-X  
DOI: <http://dx.doi.org/10.1016/j.psychres.2017.06.080>  
Reference: PSY10627

To appear in: *Psychiatry Research*

Received date: 24 October 2016  
Revised date: 23 June 2017  
Accepted date: 26 June 2017

Cite this article as: Vahid Khosravani, Farangis Sharifi Bastan, Mehdi Samim Ardestani and Razieh Jamaati Ardakani, Early maladaptive schemas and suicidal risk in an Iranian sample of patients with obsessive-compulsive disorder *Psychiatry Research*, <http://dx.doi.org/10.1016/j.psychres.2017.06.080>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and a review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain

## Early maladaptive schemas and suicidal risk in an Iranian sample of patients with obsessive-compulsive disorder

Vahid Khosravani <sup>a,\*</sup>, Farangis Sharifi Bastan <sup>b</sup>, Mehdi Samimi Ardestani <sup>c</sup>, Razieh Jamaati Ardakani <sup>d</sup>

<sup>a</sup> Clinical Research Development Center of Loghman Hakim Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

<sup>b</sup> Psychosocial Injuries Research Centre, Ilam University of Medical Sciences, Ilam, Iran.

<sup>c</sup> Departments of Psychiatry, Imam Hossein Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

<sup>d</sup> Department of Psychology, Ardakan University, Ardakan, Yazd, Iran.

\*Corresponding author. vahid.psy@gmail.com

Abstract

There are few studies on suicidal risk and its related factors in patients diagnosed with obsessive-compulsive disorder (OCD). This study investigated the associations of early maladaptive schemas, OC symptom dimensions, OCD severity, depression and anxiety with suicidality (i.e., suicidal ideation and suicide attempts) in OCD patients. Sixty OCD outpatients completed the Scale for Suicide Ideation (SSI), the Young Schema Questionnaire-Short Form (YSQ-SF), the Yale-Brown Obsessive Compulsive Scale (Y-BOCS), the Dimensional Obsessive-Compulsive Scale (DOCS) and the Depression Anxiety Stress Scales (DASS-21). 51.7% of patients had lifetime suicide attempts and 75% had suicidal ideation. OCD patients with lifetime suicide attempts exhibited significantly higher scores on early maladaptive schemas than those without such attempts. Logistic regression analysis revealed that the mistrust/abuse schema and the OC symptom dimension of unacceptable thoughts explained lifetime suicide attempts. The mistrust/abuse schema, unacceptable thoughts and depression significantly predicted suicidal ideation. These findings indicated that the mistrust/abuse schema may contribute to high suicidality in OCD patients. Also, patients suffering from unacceptable thoughts need to be assessed more carefully for warning signs of suicide.

**Keywords**

متن کامل مقاله

دریافت فوری ←

**ISI**Articles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات