

Author's Accepted Manuscript

Associations between severity of anxiety and clinical and biological features of major affective disorders

Fernanda Liboni Cavicchioli, Michael Maes, Chutima Roomruangwong, Kamila Landucci Bonifacio, Decio Sabbatini Barbosa, George Anderson, Heber Odebrecht Vargas, Sandra Odebrecht Vargas Nunes



PII: S0165-1781(17)31181-2
DOI: <https://doi.org/10.1016/j.psychres.2017.11.024>
Reference: PSY10971

To appear in: *Psychiatry Research*

Received date: 28 June 2017
Revised date: 25 October 2017
Accepted date: 6 November 2017

Cite this article as: Fernanda Liboni Cavicchioli, Michael Maes, Chutima Roomruangwong, Kamila Landucci Bonifacio, Decio Sabbatini Barbosa, George Anderson, Heber Odebrecht Vargas and Sandra Odebrecht Vargas Nunes, Associations between severity of anxiety and clinical and biological features of major affective disorders, *Psychiatry Research*, <https://doi.org/10.1016/j.psychres.2017.11.024>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Associations between severity of anxiety and clinical and biological features of major affective disorders.

Fernanda Liboni Cavicchioli^a, Michael Maes^{a,b,c,d*}, Chutima Roomruangwong^b, Kamila Landucci Bonifacio^a, Decio Sabbatini Barbosa^a, George Anderson^e, Heber Odebrecht Vargas^a, Sandra Odebrecht Vargas Nunes^a

^aHealth Sciences Graduation Program, Health Sciences Center, State University of Londrina, Brazil.

^bDepartment of Psychiatry, Chulalongkorn University, Bangkok, Thailand.

^cImpact Strategic Research Center, Deakin University, Geelong, Australia

^dDepartment of Psychiatry, Medical University of Plovdiv, Plovdiv, Bulgaria

^eCRC, Scotland and London, UK

dr.michaelmaes@hotmail.com

<http://scholar.google.co.th/citations?user=1wzMZ7UAAAAJ&hl=th&oi=ao>

*Corresponding author: Prof. Dr. Michael Maes, M.D., Ph.D. IMPACT Strategic Research Center, Barwon Health, Deakin University, Geelong, Vic, Australia

Abstract

Patients with major affective disorders (MAFD) with comorbid anxiety show a greater functional impairment than those without anxiety. The aim of this study is to delineate the associations between severity of anxiety in MAFD, namely bipolar disorder (BD) and major depression (MDD), and MAFD characteristics and serum high-density lipoprotein (HDL)-cholesterol levels. Recruited were 82 participants with anxiety disorders and 83 without

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات