Accepted Manuscript

Tau, Amyloid, and Cascading Network Failure across the Alzheimer's disease Spectrum

David T. Jones, Jonathan Graff-Radford, Val J. Lowe, Heather J. Wiste, Jeffrey L. Gunter, Matthew L. Senjem, Hugo Botha, Kejal Kantarci, Bradley F. Boeve, David S. Knopman, Ronald C. Petersen, Clifford R. Jack, Jr.

PII: S0010-9452(17)30324-6

DOI: 10.1016/j.cortex.2017.09.018

Reference: CORTEX 2139

To appear in: Cortex

Received Date: 14 March 2017 Revised Date: 2 August 2017

Accepted Date: 26 September 2017

Please cite this article as: Jones DT, Graff-Radford J, Lowe VJ, Wiste HJ, Gunter JL, Senjem ML, Botha H, Kantarci K, Boeve BF, Knopman DS, Petersen RC, Jack Jr. CR, Tau, Amyloid, and Cascading Network Failure across the Alzheimer's disease Spectrum, *CORTEX* (2017), doi: 10.1016/j.cortex.2017.09.018.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT

Tau, Amyloid, and Cascading Network Failure across the Alzheimer's disease Spectrum

David T. Jones^{1,2*}, Jonathan Graff-Radford¹, Val J. Lowe², Heather J. Wiste³, Jeffrey L. Gunter^{2,4}, Matthew L. Senjem^{2,4}, Hugo Botha¹, Kejal Kantarci², Bradley F. Boeve¹, David S. Knopman¹, Ronald C. Petersen¹, Clifford R. Jack, Jr.²

Affiliations:

¹Department of Neurology, ²Department of Radiology, ³Department of Health Sciences Research,

Contact:

*Correspondence to: David T. Jones, Mayo Clinic, 200 First Street S.W., Rochester, MN 55905 jones.david@mayo.edu

Running Title:

Tau, amyloid, and networks

Abstract:

Functionally related brain regions are selectively vulnerable to Alzheimer's disease pathophysiology. However, molecular markers of this pathophysiology (i.e., beta-amyloid and tau aggregates) have discrepant spatial and temporal patterns of progression within these selectively vulnerable brain regions. Existing reductionist pathophysiologic models cannot account for these large-scale spatiotemporal inconsistencies. Within the framework of the recently proposed cascading network failure model of Alzheimer's disease, however, these large-scale patterns are to be expected. This model postulates the following: 1) a tau-associated, circumscribed network disruption occurs in brain regions specific to a given phenotype in clinically normal individuals; 2) this disruption can trigger phenotype independent, stereotypic, and amyloid-associated compensatory brain network changes indexed by

⁴Department of Information Technology, Mayo Clinic, Rochester, MN, USA.

دريافت فورى ب متن كامل مقاله

ISIArticles مرجع مقالات تخصصی ایران

- ✔ امكان دانلود نسخه تمام متن مقالات انگليسي
 - ✓ امكان دانلود نسخه ترجمه شده مقالات
 - ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
 - ✓ امكان دانلود رايگان ۲ صفحه اول هر مقاله
 - ✔ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
 - ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات