

Accepted Manuscript

Title: Aluminium in brain tissue in autism

Authors: Matthew Mold, Dorcas Umar, Andrew King, Christopher Exley


PII: S0946-672X(17)30876-3
DOI: <https://doi.org/10.1016/j.jtemb.2017.11.012>
Reference: JTEMB 26015

To appear in:

Received date: 26-10-2017
Revised date: 21-11-2017
Accepted date: 23-11-2017

Please cite this article as: Mold Matthew, Umar Dorcas, King Andrew, Exley Christopher. Aluminium in brain tissue in autism. *Journal of Trace Elements in Medicine and Biology* <https://doi.org/10.1016/j.jtemb.2017.11.012>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Aluminium in brain tissue in autism

Matthew Mold^a, Dorcas Umar^b, Andrew King^c, Christopher Exley^{a*}

^aThe Birchall Centre, Lennard-Jones Laboratories, Keele University, Staffordshire, ST5 5BG, United Kingdom.

^bLife Sciences, Keele University, Staffordshire, ST5 5BG, United Kingdom.

^cDepartment of Clinical Neuropathology, Kings College Hospital, London, SE5 9RS, United Kingdom.

ABSTRACT

Autism spectrum disorder is a neurodevelopmental disorder of unknown aetiology. It is suggested to involve both genetic susceptibility and environmental factors including in the latter environmental toxins. Human exposure to the environmental toxin aluminium has been linked, if tentatively, to autism spectrum disorder. Herein we have used transversely heated graphite furnace atomic absorption spectrometry to measure, for the first time, the aluminium content of brain tissue from donors with a diagnosis of autism. We have also used an aluminium-selective fluor to identify aluminium in brain tissue using fluorescence microscopy. The aluminium content of brain tissue in autism was consistently high. The mean (standard deviation) aluminium content across all 5 individuals for each lobe were 3.82(5.42), 2.30(2.00), 2.79(4.05) and 3.82(5.17) $\mu\text{g/g}$ dry wt. for the occipital, frontal, temporal and parietal lobes respectively. These are some of the highest values for aluminium in human brain tissue yet recorded and one has to question why, for example, the aluminium content of the occipital lobe of a 15 year old boy would be 8.74 (11.59) $\mu\text{g/g}$ dry wt.?

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات