

Author's Accepted Manuscript

THE ASSOCIATION OF SERUM NESFATIN-1 AND GHRELIN LEVELS WITH METABOLIC SYNDROME IN SCHIZOPHRENIA PATIENTS

Kubranur Unal, Rabia Nazik Yuksel, Turan
Turhan, Sevilay Sezer, Elif Tatlidil Yaylaci



PII: S0165-1781(17)30964-2
DOI: <https://doi.org/10.1016/j.psychres.2017.12.041>
Reference: PSY11077

To appear in: *Psychiatry Research*

Received date: 30 May 2017
Revised date: 5 November 2017
Accepted date: 14 December 2017

Cite this article as: Kubranur Unal, Rabia Nazik Yuksel, Turan Turhan, Sevilay Sezer and Elif Tatlidil Yaylaci, THE ASSOCIATION OF SERUM NESFATIN-1 AND GHRELIN LEVELS WITH METABOLIC SYNDROME IN SCHIZOPHRENIA PATIENTS, *Psychiatry Research*, <https://doi.org/10.1016/j.psychres.2017.12.041>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

THE ASSOCIATION OF SERUM NESFATIN-1 AND GHRELIN LEVELS WITH METABOLIC SYNDROME IN SCHIZOPHRENIA PATIENTS

Kubranur Unal^a, Rabia Nazik Yüksel^{b*}, Turan Turhan^c, Sevilay Sezer^c, Elif Tatlıdil Yaylacı^b

^aDepartment Of Biochemistry, Polatlı State Hospital, Ankara, Turkey

^bDepartment Of Psychiatry, Ankara Numune Training And Research Hospital, Ankara, Turkey

^cDepartment Of Biochemistry, Ankara Numune Training And Research Hospital, Ankara, Turkey

*Corresponding Author: Rabia Nazik Yüksel. Address: Talatpasa Bulvarı No:5 Ankara Numune Eğitim ve Araştırma Hastanesi, Altındag, Ankara. Phone Number: +903125085612. rabianazik@gmail.com

Abstract

Nesfatin-1 and ghrelin are two hormones which has opposite effects and play role in food intake. This study was planned on the idea that both metabolic syndrome and psychiatric disorders are associated with nesfatin-1 and ghrelin. In this study, it was aimed to investigate the levels of ghrelin and nesfatin-1 in patients with schizophrenia, by taking confounding factor as the metabolic syndrome (MS). 55 patients with schizophrenia and 33 healthy controls were included in the study. 11 out of the 55 patients (%20) has MS. Serum ghrelin and nesfatin-1 levels of schizophrenia patients with MS have been compared with both healthy controls and schizophrenia patients without MS. Patients with schizophrenia had significantly higher serum nesfatin-1 levels compared to healthy controls. But serum ghrelin levels was not different in both groups. Serum nesfatin-1 concentrations were significantly higher in the schizophrenia patients with MS (10.51-350.8 pg/ml) with respect to the healthy control group (4.86-68.91 pg/ml). There was no significant statistical difference between the three groups in terms of ghrelin levels. Our findings suggests that, MS presence also contributed to significantly high levels of nesfatin-1 level. Nesfatin-1 may have a part in a novel studies regarding the treatment of schizophrenia and its metabolic effects.

Keywords: Schizophrenia, Nesfatin-1, Ghrelin, Metabolic Syndrome, Schizophrenia treatment

1. Introduction

Low activity, unbalanced nourishment, cigarettes and antipsychotic drugs used in schizophrenia patients would cause weight gain, diabetes, and impairments in their lipid profile (Kaiya et al., 1989; McIntyre et al., 2001; Henderson, 2002; Association, 2004). Compared to the general population, abdominal obesity, glucose intolerance, or diabetes mellitus, dyslipidemia, hypertension, and cardiovascular diseases occur more frequently in schizophrenia patients. These risk factors are called metabolic syndrome, which increases the cardiovascular disease risks by 25-50% (García-Bueno et al., 2014). Life expectancy of schizophrenia patients are shorter when compared to that of the general population. Suicide and cardiovascular disease are the most common causes of death in schizophrenia patients (Kaiya et al., 1989). It was found that the basis of metabolic syndrome is a damaged response to the tissues in insulin and insulin resistance dependent hyperinsulinemia is developed (Wilcox, 2005)

It was hypothesized that neuroendocrine molecules play a role in the etiology of metabolic syndrome (Kaiya et al., 1989; Das and Khan, 1998). In recent studies, the important effects of molecules such as ghrelin and nesfatin-1 on weight and hunger mechanisms were reported (Gervois et al., 2007; Saetre et al., 2007). Ghrelin is a lipopeptide

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات