

## Accepted Manuscript

Title: Burnout risk profiles among French psychologists

Authors: Sophie Berjot, Emin Altintas, François-Xavier Lesage, Elisabeth Grebot

PII: S2213-0586(17)30018-9  
DOI: <https://doi.org/10.1016/j.burn.2017.10.001>  
Reference: BURN 56



To appear in:

Received date: 10-4-2017  
Revised date: 21-9-2017  
Accepted date: 1-10-2017

Please cite this article as: { <https://doi.org/>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## **Burnout risk profiles among French psychologists**

Sophie Berjot,<sup>1</sup> Emin Altintas,<sup>2</sup> François-Xavier Lesage,<sup>3</sup>

Elisabeth Grebot,<sup>3</sup>

1. Laboratoire Cognition, Santé & Socialisation C2S  
Université de Reims Champagne-Ardenne - France

2. Laboratoire PSITEC  
Université Lille Nord de France – France

3. Laboratoire EPSYLON, CHU de Montpellier - France

4. Laboratoire de psychopathologie et des processus de santé  
Université Paris Descartes, Sorbonne Paris Cité - France

Address correspondence to Sophie Berjot, Laboratoire Cognition, Santé & Socialisation, UFR Lettres et Sciences Humaines, Université de Reims Champagne-Ardenne, 57, rue Pierre-Taittinger, 51096 Reims Cedex – France or e-mail ([sophie.berjot@univ-reims.fr](mailto:sophie.berjot@univ-reims.fr))

We would like to thank the IRIST (Institut de Recherche et d'Intervention en Santé au Travail - Institute of Research and Intervention in Health at Work) for its help in carrying this study

Key words: Burnout, psychologists, prevalence

Word count: 3 711

### **Highlights**

- The use of cut-offs scores to assess burnout's prevalence may be misleading
- The use of cluster analysis to classify workers into risk groups may represent an alternative
- This type of analysis can help to highlight groups at risk of burnout, and different risk constellations within a profession.
- There are different patterns of burnout according to age, working place, and type of contract.

متن کامل مقاله

دریافت فوری ←

**ISI**Articles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات