
Author's Accepted Manuscript

Treatment Outcomes for Inpatients with Obsessive-Compulsive Personality Disorder: An Open Comparison Trial

Ryan Smith, Christopher Shepard, Anika Wiltgen, Katrina Rufino, J. Christopher Fowler



 PII:
 S0165-0327(16)31056-4

 DOI:
 http://dx.doi.org/10.1016/j.jad.2016.12.002

 Reference:
 JAD8663

To appear in: Journal of Affective Disorders

Received date: 22 June 2016 Revised date: 22 November 2016 Accepted date: 4 December 2016

Cite this article as: Ryan Smith, Christopher Shepard, Anika Wiltgen, Katrina Rufino and J. Christopher Fowler, Treatment Outcomes for Inpatients witl Obsessive-Compulsive Personality Disorder: An Open Comparison Trial *Journal of Affective Disorders*, http://dx.doi.org/10.1016/j.jad.2016.12.002

This is a PDF file of an unedited manuscript that has been accepted fo publication. As a service to our customers we are providing this early version o the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain

ACCEPTED MANUSCRIPT

Treatment Outcomes for Inpatients with Obsessive-Compulsive Personality Disorder: An

Open Comparison Trial

Ryan Smith^{a*}, Christopher Shepard^a, Anika Wiltgen^a, Katrina Rufino^{a,b}, J. Christopher Fowler^{a,c}

^aMenninger Clinic

^bUniversity of Houston Downtown

^cBaylor College of Medicine

*Corresponding Author: The Menninger Clinic, 12301 Main St., Houston, TX 77035 Tel: +1-713-275-5486, Email Address: rpsmith@menninger.edu

Abstract

Background:

The current case-control study compared rates of clinically significant and reliable change in psychopathology and global functioning, prevalence of clinical deterioration, and rates of symptom remission among adult patients with obsessive compulsive personality disorder OCPD (n=52) and well-matched inpatients with any other personality disorder (n=56) and no personality disorder (n=53).

Methods:

Propensity score matching (PSM) was utilized to select patients matched on specific criteria present in the OCPD group. Multivariate analysis of variance models measured differences in

دريافت فورى 🛶 متن كامل مقاله

- امکان دانلود نسخه تمام متن مقالات انگلیسی
 امکان دانلود نسخه ترجمه شده مقالات
 پذیرش سفارش ترجمه تخصصی
 امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
 امکان دانلود رایگان ۲ صفحه اول هر مقاله
 امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
 دانلود فوری مقاله پس از پرداخت آنلاین
 پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات
- ISIArticles مرجع مقالات تخصصی ایران