

Accepted Manuscript

Objective Assessment of Acne

Markus Becker MD, Thomas Wild MD, Christos C Zouboulis MD, PhD

PII: S0738-081X(16)30267-X
DOI: doi: [10.1016/j.clindermatol.2016.10.006](https://doi.org/10.1016/j.clindermatol.2016.10.006)
Reference: CID 7104

To appear in: *Clinics in Dermatology*



Please cite this article as: Becker Markus, Wild Thomas, Zouboulis Christos C, Objective Assessment of Acne, *Clinics in Dermatology* (2016), doi: [10.1016/j.clindermatol.2016.10.006](https://doi.org/10.1016/j.clindermatol.2016.10.006)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Objective Assessment of Acne

Authors and affiliations

Markus Becker, MD¹, Thomas Wild, MD¹, Christos C Zouboulis, MD, PhD¹

¹ Departments of Dermatology, Venereology, Allergology and Immunology, Dessau Medical Center, Auenweg 38, 06847 Dessau, Germany

Corresponding author

Markus Becker¹

¹ Departments of Dermatology, Venereology, Allergology and Immunology, Dessau Medical Center, Auenweg 38, 06847 Dessau, Germany

Emails: Markus Becker: markus.becker@klinikum-dessau.de

Thomas Wild: thomas.wild@klinikum-dessau.de

Christos C. Zouboulis: christos.zouboulis@klinikum-dessau.de

Abstract

A precise and reliable assessment of acne severity is unarguably the most essential clinical method, when it comes to monitoring and choosing optimal treatment in the daily practice. Since the early 1960's, different severity assessment systems have been described in the literature. The two commonly used concepts are global gradings and lesion counting. Both systems have been controversially discussed as to which is more reliable and providing an objective outcome measurement tool; however, both have some subjectivity involved.. More objective methods for assessing the severity of acne vulgaris include photography, fluorescence photography, polarized light photography, video microscopy, and multispectral imaging; Such techniques have limitations involving high cost, complex and sophisticated apparatuses, and a sometimes time consuming imaging process. There are newly developed technologies that could avoid the problems of inter- and intra-rater subjectivity.

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات