

## Accepted Manuscript

Title: Prolonged striatal disinhibition as a chronic animal model of tic disorders

Authors: Esther Vinner, Michal Israelashvili, Izhar Bar-Gad

PII: S0165-0270(17)30060-2  
DOI: <http://dx.doi.org/doi:10.1016/j.jneumeth.2017.03.003>  
Reference: NSM 7689

To appear in: *Journal of Neuroscience Methods*

Received date: 27-9-2016  
Revised date: 20-1-2017  
Accepted date: 3-3-2017

Please cite this article as: Vinner Esther, Israelashvili Michal, Bar-Gad Izhar. Prolonged striatal disinhibition as a chronic animal model of tic disorders. *Journal of Neuroscience Methods* <http://dx.doi.org/10.1016/j.jneumeth.2017.03.003>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



**Prolonged striatal disinhibition as a chronic animal model of tic disorders**

Esther Vinner, Michal Israelashvili, Izhar Bar-Gad\*

*Gonda Multidisciplinary Brain Research Center, Bar Ilan University, Ramat Gan, Israel*

**Highlights**

- The chronic model is an adaptation of the acute striatal disinhibition model.
- Mini osmotic pumps were used for prolonged bicuculline infusion to the striatum.
- Motor tics were expressed throughout the infusion period (multiple days).
- Tics were accompanied by stereotypical LFP spikes throughout the period.
- Neuronal activity around the tic time was stereotypic and stable across days.

**Abstract***Background*

Experimental findings and theoretical models have associated Tourette syndrome with abnormal striatal inhibition. The expression of tics, the hallmark symptom of this disorder, has been transiently induced in non-human primates and rodents by the injection of GABA<sub>A</sub> antagonists into the striatum, leading to temporary disinhibition.

*New method*

The novel chronic model of tic expression utilizes mini-osmotic pumps implanted subcutaneously in the rat back for prolonged infusion of bicuculline into the dorsolateral striatum.

*Results*

Tics were expressed on the contralateral side to the infusion over a period of multiple days. Tic expression was stable, and maintained similar properties throughout the infusion period. Electrophysiological recordings revealed the existence of tic-related local field potential spikes and individual neuron activity changes that remained stable throughout the infusion period.

*Comparison with existing methods*

The striatal disinhibition model provides a unique combination of face validity (tic expression) and construct validity (abnormal striatal inhibition) but is limited to sub-hour periods. The new chronic model extends the period of tic expression to multiple days and thus enables the study of tic dynamics and the effects of behavior and pharmacological agents on tic expression.

*Conclusions*

متن کامل مقاله

دریافت فوری ←

**ISI**Articles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات