

Author's Accepted Manuscript

Alcohol use disorders are associated with increased affective lability in bipolar disorder

Trine Vik Lagerberg, Sofie Ragnhild Aminoff, Monica Aas, Thomas Bjella, Chantal Henry, Marion Leboyer, Geir Pedersen, Frank Bellivier, Romain Icick, Ole A. Andreassen, Bruno Etain, Ingrid Melle



PII: S0165-0327(16)30690-5
DOI: <http://dx.doi.org/10.1016/j.jad.2016.09.062>
Reference: JAD8536

To appear in: *Journal of Affective Disorders*

Received date: 26 April 2016
Revised date: 15 August 2016
Accepted date: 5 September 2016

Cite this article as: Trine Vik Lagerberg, Sofie Ragnhild Aminoff, Monica Aas, Thomas Bjella, Chantal Henry, Marion Leboyer, Geir Pedersen, Frank Bellivier, Romain Icick, Ole A. Andreassen, Bruno Etain and Ingrid Melle, Alcohol use disorders are associated with increased affective lability in bipolar disorder *Journal of Affective Disorders*, <http://dx.doi.org/10.1016/j.jad.2016.09.062>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain

Alcohol use disorders are associated with increased affective lability in bipolar disorder

Trine Vik Lagerberg^{A*}, Sofie Ragnhild Aminoff^{A,B}, Monica Aas^A, Thomas Bjella^A, Chantal Henry^{C,D,E,F}, Marion Leboyer^{C,D,F,G}, Geir Pedersen^{A,H}, Frank Bellivier^{F,I,J,K,L}, Romain Icick^{I,J,K,L}, Ole A. Andreassen^{A,G}, Bruno Etain^{F,I,J,K,L}, Ingrid Melle^A.

- A) NORMENT, KG Jebsen Centre for Psychosis Research, Oslo University Hospital and Institute of Clinical Medicine, University of Oslo, Oslo, Norway.
- B) Department of Specialized Inpatient Treatment, Division of Mental Health Services, Akershus University Hospital, Norway
- C) AP-HP, Hôpital H. Mondor – A. Chenevier, DHU Pepsy, Pôle de Psychiatrie, Créteil 94000, France
- D) Inserm, U955, Créteil 94000, France
- E) Institut Pasteur, Unité Perception et Mémoire, F-75015 Paris, France
- F) ENBREC, European Network of Bipolar Research Expert Centres, Paris, France
- G) Fondation FondaMental, Créteil, France
- H) Oslo University Hospital, Department of Personality Psychiatry, Division of Mental Health and Addiction, Oslo, Norway
- I) AP-HP, GH Saint-Louis – Lariboisière – F. Widal, Pôle de Psychiatrie et de Médecine Addictologique, 75475 Paris cedex 10, France
- J) Inserm, U1144, Paris, F-75006, France
- K) Université Paris Descartes, UMR-S 1144, Paris, F-75006, France
- L) Université Paris Diderot, Sorbonne Paris Cité, UMR-S 1144, Paris, F-75013, France

*Corresponding author. Mail: t.v.lagerberg@medisin.uio.no, phone: +47 99020335

ABSTRACT

Background: Affective dysregulation is a core feature of bipolar disorder (BD), and inter-episodic affect lability is associated with more severe outcomes including comorbidity. Rates of daily tobacco smoking and substance use disorders in BD are high. Knowledge regarding relationships between affective lability and abuse of the most commonly used substances such as tobacco, alcohol and cannabis in BD is limited.

Methods: We investigated whether dimensions of inter-episodic affective lability as measured with the Affective Lability Scale – short form (ALS-SF) were associated with lifetime daily tobacco use or

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات