Accepted Manuscript

Preparation of molecularly imprinted electrochemical sensor for lphenylalanine detection and its application

Nihal Ermiş, Lokman Uzun, Adil Denizli



Please cite this article as: Nihal Ermiş, Lokman Uzun, Adil Denizli, Preparation of molecularly imprinted electrochemical sensor for l-phenylalanine detection and its application. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Jeac(2017), doi:10.1016/j.jelechem.2017.11.025

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT

Preparation of Molecularly Imprinted Electrochemical Sensor For L-Phenylalanine Detection and its Application

Nihal Ermiş^a,^{*} Lokman Uzun^b, Adil Denizli^b

^aOndokuz Mayıs University, Faculty of Science and Letters, Department of Chemistry, Samsun, Turkey ^bHacettepe University, Faculty of Science, Department of Chemistry, Ankara, Turkey Corresponding Author : nihal.ermis@omu.edu.tr

Abstract

For selective and sensitive determination of L-phenylalanine (L-Phe) a novel and reliable electrochemical sensor has been developed based on molecular imprinting technique. Thiophen-3-carbonyl tryptophan (TP3C-Trp) has been synthesized and used as a novel monomer. L-Phe was used as template molecule and imprinted in presence of thiophen and thiophen-3-carbonyl tryptophan (TP3C-Trp), used as functional monomers. As a comparison a non-modified polymer surface was also prepared under same circumstances without template molecule, L-Phe. The characterization studies of both non-imprinted (NIP) and L-Phenylalanine imprinted (MIP) electrochemical sensor surfaces were made by using atomic force microscopy (AFM), Fourier transform infrared spectroscopy (FTIR), ellipsometry and contact angle measurements. The electrochemical characterization analysis were made via square wave and cyclic voltammetry. The linearity range and the detection limit of the developed sensor were obtained as 1.0x10⁻⁸–1.0x10⁻⁷ M and 1.0x10⁻⁹ M, respectively.

1. Introduction

Phenylalanine (Phe) is an essential non-polar amino acid and shows hydrophobic feature because of its aromatic group. It is also metabolic precursor of non-essential amino acid tyrosine and derivatives. It is used in pharmaceutical industry, to fabricate some antibiotics like bacitracin, gramicidin, tirocidin and in food industry used as sweetener, aspartame [1,2]. It is converted to tyrosine (Tyr) in presence of phenylalanine hydroxylase enzyme (PAH). Tyrosine is the precursor of monoamine neurotransmitters, such as dopamine, epinephrine and norepinephrine which are associated with depression and lack of motivation. Therefore Phe is important to maintain adequate levels of monoamine neurotransmitters in the brain. When L-phenylalanine crosses into the brain in the presence of PAH, it is converted to L-tyrosine and then to L-dopa which is further converted into the three neurotransmitters mentioned above. Adrenaline and noradrenaline are synthesised from dopamine, which is itself made from the amino acid phenylalanine via tyrosine. It is well known that

دريافت فورى 🛶 متن كامل مقاله

- امکان دانلود نسخه تمام متن مقالات انگلیسی
 امکان دانلود نسخه ترجمه شده مقالات
 پذیرش سفارش ترجمه تخصصی
 امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
 امکان دانلود رایگان ۲ صفحه اول هر مقاله
 امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
 دانلود فوری مقاله پس از پرداخت آنلاین
 پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات
- ISIArticles مرجع مقالات تخصصی ایران