



Temporal stability of multiple response systems to 7.5% carbon dioxide challenge



Roxann Roberson-Nay^{a,*}, Eugenia I. Gorlin^b, Jessica R. Beadel^b, Therese Cash^a, Scott Vrana^a, Bethany A. Teachman^b

^a Virginia Commonwealth University, United States

^b University of Virginia, United States

ARTICLE INFO

Article history:

Received 6 April 2016

Received in revised form 18 January 2017

Accepted 31 January 2017

Available online 2 February 2017

Keywords:

Carbon dioxide hypersensitivity

Panic risk

Temporal stability

Reliability

Subjective anxiety

Respiratory rate

Heart rate

Skin conductance

Panic attack

Panic symptoms

ABSTRACT

Self-reported anxiety, and potentially physiological response, to maintained inhalation of carbon dioxide (CO₂) enriched air shows promise as a putative marker of panic reactivity and vulnerability. Temporal stability of response systems during low-dose, steady-state CO₂ breathing challenge is lacking. Outcomes on multiple levels were measured two times, one week apart, in 93 individuals. Stability was highest during the CO₂ breathing phase compared to pre-CO₂ and recovery phases, with anxiety ratings, respiratory rate, skin conductance level, and heart rate demonstrating good to excellent temporal stability (ICCs \geq 0.71). Cognitive symptoms tied to panic were somewhat less stable (ICC = 0.58) than physical symptoms (ICC = 0.74) during CO₂ breathing. Escape/avoidance behaviors and DSM-5 panic attacks were not stable. Large effect sizes between task phases also were observed. Overall, results suggest good–excellent levels of temporal stability for multiple outcomes during respiratory stimulation via 7.5% CO₂.

© 2017 Elsevier B.V. All rights reserved.

Carbon dioxide (CO₂) hypersensitivity has been studied among persons with panic disorder and other psychiatric conditions, as well as in the general population, and appears to represent a fairly well validated measure of panic reactivity and vulnerability (Battaglia et al., 2007; Coryell, 1997; Coryell, Fyer, Pine, Martinez, & Arndt, 2001; Griez, de Loof, Pols, Zandbergen, & Lousberg, 1990; Perna, Cocchi, Bertani, Arancio, & Bellodi, 1995; Perna, Cocchi, Allevi, Bussi, & Bellodi, 1999). Studies using CO₂ enriched air have relied on a variety of CO₂ air concentrations, generally ranging from 5% to 35% CO₂. Of the concentrations used in clinical research, 35% CO₂ is the most commonly used CO₂ challenge intensity, likely owing to its procedural ease, wherein participants inhale one or two vital capacity breaths and their anxiety and symptomatic response pre- and post-inhalation are assessed (Amaral et al., 2013). By contrast, the lower CO₂ concentrations (e.g., 5%, 7.5%) allow researchers to continuously measure autonomic levels

(e.g., respiratory rate, heart rate) as well as self-report of anxiety and symptomatic response, permitting the modeling of change over time.

The National Institute for Mental Health (NIMH) recently launched the Research Domain Criteria (RDoC) project as part of their strategic plan to develop novel ways of classifying psychiatric disorders based on dimensions of observable behaviors and brain functions (Cuthbert & Insel, 2010, 2013; Insel & Cuthbert, 2009; Insel et al., 2010; Sanislow et al., 2010). RDoC aims to serve as a framework for new approaches to research on mental disorders using fundamental dimensions that cut across traditional disorder categories. It is hoped that these fundamental dimensions will closely align with mechanisms that underlie psychopathology at various biological and behavioral levels. The CO₂ challenge fits well within the RDoC matrix as a “paradigm” under the Negative Valence Systems and provides for a number of units of analysis, including (but not limited to) physiology, behavior, neural circuits, and self-report.

Given this conceptual shift from diagnostic level phenotypes to dimensional construct measures that fall under higher order diagnoses, the need for psychometrically substantiated measures is crucial. Moreover, given that sensitivity to CO₂ is hypothesized to

* Corresponding author at: Department of Psychiatry, Virginia Commonwealth University, Virginia Institute for Psychiatric and Behavioral Genetics, P.O. Box 980489, Richmond, VA 23298, United States.

E-mail address: roxann.robersonnay@vcuhealth.org (R. Roberson-Nay).

be a biologically based trait marker of panic vulnerability (Battaglia et al., 2007; Coryell, 1997; Coryell et al., 2001; Griez et al., 1990; Perna et al., 1995, 1999), it is important to determine whether this marker is consistently expressed across time. Two previous studies examined the test-retest reliability of self-reported anxiety and symptom response to inhalation of 35% CO₂, but these studies relied on a small, clinical sample (Verburg, de Leeuw, Pols, & Griez, 1997) and a high- versus low-risk sample (Coryell & Arndt, 1999). Both studies documented reasonable reliability of symptomatic response, particularly smothering sensations, faintness, and dizziness, as well as greater reliability for anxiety rated post versus prior to 35% CO₂ inhalation. Physiological and behavioral markers were not measured. Using a within-session repeated measures design, participants' cardiac, electrodermal, and self-report of anxiety were measured in response to eight, 20 s inhalations of 20% CO₂ enriched air (Forsyth, Eifert, & Canna, 2000). Means testing indicated no significant trial effects, reflecting stability of measures within session. Collectively these studies suggest satisfactory stability of physiology and subjective response systems to high dose CO₂ enriched air. Thus, although the stability of anxiety, symptom, and physiological response to high dose CO₂ has been established, a temporal stability study of low dose, steady-state CO₂ breathing has not been undertaken.

The objective of the current study is to determine the test-retest reliability of self-reported anxiety and panic symptom response as well as behavioral and physiological markers during the steady-state 7.5% CO₂ challenge across two time points. Study outcomes will inform future research and clinical assessments where stable markers are needed. Moderate levels of stability of subjective and symptom report is expected based on previous studies examining stability of response to higher dose CO₂ concentrations (Coryell & Arndt, 1999; Forsyth et al., 2000; Verburg et al., 1997). Extant studies of physiological reliability suggest reasonably good temporal consistency (Forsyth et al., 2000; Schmidt et al., 2002). Thus, satisfactory reliability estimates also are expected for physiological outcomes.

1. Methods

1.1. Participants

Participants were recruited as part of a larger study of CO₂ hypersensitivity (n=376) (Roberson-Nay, Beadel, Gorlin, Latendresse, & Teachman, 2015) at two universities from either the psychology department participant pool or recruitment fliers posted on the campus. Table 1 provides demographic and panic related characteristics for the full test-retest reliability sample (n=93) as well as for each site. A power calculation conducted before the study's start, indicated a sample size of 86 participants for reliability analysis (see Methods section for full calculation). Thus, the first, consecutive 146 participants completing Session 1

of the CO₂ hypersensitivity study at Sites 1 and 2 were invited back to participate in Session 2 of the reliability study. Of the 146 participants invited back for Session 2, 93 (64%) agreed to return. We, therefore, slightly exceeded our power calculation estimation of 86 individuals for reliability analysis.

Study participants at both sites were compensated financially or with course credit; the majority (89%) received course credit. Site 2 recruited participants based on a participant's Anxiety Sensitivity Index (ASI; Reiss, Peterson, Gursky, & McNally, 1986) score, which was completed during a pre-screening session. Participants were recruited based on ASI scores to ensure variance on this important panic and CO₂ response risk factor (Blechert, Wilhelm, Meuret, Wilhelm, & Roth, 2013; Korte & Schmidt, 2012; Schmidt, Lerew, & Jackson, 1997, Schmidt, Lerew, & Jackson, 1999; Schmidt, Zvolensky, & Maner, 2006). Thus, Site 2 sent recruitment e-mails to approximately equivalent numbers of students scoring within each quartile of the distribution of ASI scores (Peterson & Reiss, 1992). No pre-screen ASI selection criteria was used for participants who participated for financial compensation or who participated at Site 1; Site 1 was unable to include a pre-screening ASI screen given differences in the participant pool infrastructure. Nonetheless, ASI score distributions and means/variance did not differ between sites (see Table 1). ASI scores also did not differ between participants participating for course credit versus financial compensation ($t(1.89)=0.95, p=0.34$, Cohen's $d=0.33$).

Of the 146 participants invited to return for session 2, there were no differences between participants who did and did not return to participate in Session 2 on the following measures, which were assessed at Session 1: repeated measurements of anxiety ratings, Diagnostic Symptom Questionnaire scores, and all physiological measures assessed during the three phases of the CO₂ challenge, as well as the Anxiety Sensitivity Index (all p 's > 0.05). The one exception was a non-significant trend ($t(1142)=0.85, p=0.40$, Cohen's $d=0.15$) for the anxiety rating measured during the attachment of the facemask while breathing room air; participants who returned to partake in Session 2 had a slightly higher mean SUDS rating ($M=23.6$) compared to participants who did not return ($M=19.5$). There also was no difference in the number of females/males ($\chi^2=0.012, p=0.91$) who did and did not return for the Session 2 assessment. Finally, rate of early termination of the CO₂ challenge at Session 1 was nearly identical for those participants who agreed to return (21.5%) versus did not agree to return (21.6%) to participate in Session 2 ($\chi^2=0.00, p=0.99$). These results suggest that there were not significant differences between participants who did versus did not return to participate in the Session 2 CO₂ challenge on primary study outcomes.

To maintain consistency across the test and retest sessions, all Session 2 visits were conducted in the same room as Session 1. Although efforts were made to have the same study team member complete Session 1 and Session 2 with the same test-retest reliability participant, this was not always possible. In total, however,

Table 1
Demographic and panic related characteristics measured before 7.5% CO₂ challenge for the full reliability sample and by study site.

	Full Sample n=93	Site 1 n=52	Site 2 n=41	t/χ^2	p
Age	19.9 (4.2)	20.3 (5.2)	19.3 (2.3)	1.1	0.29
Sex, % female	52.7	62.7	48.6	1.7	0.19
Self-reported Race, %					
African American	22.0	30.0	9.1	5.5	0.14
Asian	14.6	12.2	18.2		
Caucasian	57.3	51.0	66.7		
Other/More than one Race	6.1	6.1	6.1		
Self-reported Ethnicity, % Hispanic	12.8	9.8	17.1	1.0	0.32
PDSS, % above screening cut-off score ≥ 8	14.0	17.6	8.6	0.94	0.23
ASI Total Score	18.8 (10.2)	18.7 (9.1)	18.9	-0.05	0.96

PDSS = Panic Disorder Severity Scale; ASI = Anxiety Sensitivity Index.

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات