

## Author's Accepted Manuscript

A brief cognitive therapy intervention for internalised stigma in acute inpatients who experience psychosis: a feasibility randomised controlled trial

Lisa Wood, Rory Byrne, Gabriela Enache, Tony Morrison



PII: S0165-1781(17)30554-1  
DOI: <https://doi.org/10.1016/j.psychres.2017.12.030>  
Reference: PSY11066

To appear in: *Psychiatry Research*

Received date: 29 March 2017  
Revised date: 11 December 2017  
Accepted date: 12 December 2017

Cite this article as: Lisa Wood, Rory Byrne, Gabriela Enache and Tony Morrison, A brief cognitive therapy intervention for internalised stigma in acute inpatients who experience psychosis: a feasibility randomised controlled trial, *Psychiatry Research*, <https://doi.org/10.1016/j.psychres.2017.12.030>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**A brief cognitive therapy intervention for internalised stigma in acute inpatients who experience psychosis: a feasibility randomised controlled trial**

**Lisa Wood**

**Rory Byrne**

**Gabriela Enache**

**Tony Morrison**

**Abstract**

Internalised stigma is problematic for people who experience psychosis therefore psychological interventions are required. This study examine the feasibility and acceptability of a brief Cognitive Behavioural Therapy (CBT) intervention for internalised stigma with psychiatric inpatients experiencing psychosis. A feasibility randomised controlled trial was conducted, comparing CBT with a psychoeducational (PE) control arm. Thirty participants (aged 18-65, with psychosis, and currently admitted to a psychiatric hospital) were randomised to one of two conditions. Participants were assessed at baseline, post-intervention (two weeks) and at follow-up (one month). Both interventions incorporated two hours of sessions over a two week period. The outcomes examined were internalised stigma (primary outcome), stigma, attitudes toward mental health problems, personal recovery, depression and self-esteem. Recruitment was conducted over a seven month period from five psychiatric wards. Forty five potential participants were approached and 30 (66%) consented to take part. Fifteen participants were randomised to CBT and 15 to PE. Feasibility data demonstrated that both the research process and interventions were feasible and acceptable. Examination of outcomes demonstrated that there was no identified benefit of one intervention type over

متن کامل مقاله

دریافت فوری ←

**ISI**Articles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات