


# Accepted Manuscript

Anterior transthoracic surgery with motor evoked potential monitoring for high-risk thoracic disc herniations: technique and results

Erwin Cornips, Jeroen Habets, Vivianne van Kranen - Mastenbroek, Herman Bos, Paul Bergs, Alida Postma



PII: S1878-8750(17)30875-6

DOI: [10.1016/j.wneu.2017.05.173](https://doi.org/10.1016/j.wneu.2017.05.173)

Reference: WNEU 5867

To appear in: *World Neurosurgery*

Received Date: 24 December 2016

Revised Date: 29 May 2017

Accepted Date: 30 May 2017

Please cite this article as: Cornips E, Habets J, van Kranen - Mastenbroek V, Bos H, Bergs P, Postma A, Anterior transthoracic surgery with motor evoked potential monitoring for high-risk thoracic disc herniations: technique and results, *World Neurosurgery* (2017), doi: 10.1016/j.wneu.2017.05.173.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Anterior transthoracic surgery with motor evoked potential monitoring for high-risk thoracic disc herniations: technique and results**

Erwin Cornips<sup>a</sup>, Jeroen Habets<sup>a</sup>, Vivianne van Kranen - Mastenbroek<sup>b</sup>, Herman Bos<sup>c</sup>, Paul Bergs<sup>b</sup>, Alida Postma<sup>d</sup>

<sup>a</sup>Department of Neurosurgery, Maastricht University Medical Center, Maastricht, The Netherlands

<sup>b</sup>Department of Clinical Neurophysiology, Maastricht University Medical Center, Maastricht, The Netherlands

<sup>c</sup>Department of Anesthesiology, Maastricht University Medical Center, Maastricht, The Netherlands

<sup>d</sup>Department of Radiology, Maastricht University Medical Center, Maastricht, The Netherlands

**Abstract**

*Objective*

We present our experience with anterior transthoracic surgery for high-risk thoracic disc herniations (TDHs) using motor evoked potential monitoring (MEPm).

*Methods*

We estimated surgical risk based on clinical and radiological characteristics, including antero-posterior (AP) spinal canal occupation (SCO), AP spinal cord compression (SCC), residual AP spinal cord diameter (RSCD), and intramedullary signal changes (ISC). We analyzed Anand, ASIA, and Nurick scores, anesthesiological, neurophysiological, and surgical reports. We considered >50% MEP signal deterioration significant.

*Results*

Out of 435 anterior transthoracic procedures, 77 concerned high-risk TDH(s): 69 single-level (SL), 8 multi-level (ML), 61 with clinical myelopathy, 6 with merely ISC. Mean SCO and SCC were 55.4% and 54.0% for SL, 31.8% and 33.9% for ML cases. ISC were present in 64% of SL and 50% of ML cases, calcifications in 87% of SL and 84% of ML TDHs. We initially performed 23 mini-transthoracic approaches (mini-TTAs) and 24 thoracoscopic

متن کامل مقاله

دریافت فوری ←

**ISI**Articles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات