Accepted Manuscript

Representation of steady-state visual evoked potentials elicited by luminance flicker in human occipital cortex: An electrocorticography study

Benjamin Wittevrongel, Elvira Khachatryan, Mansoureh Fahimi Hnazaee, Evelien Carrette, Leen De Taeye, Alfred Meurs, Paul Boon, Dirk Van Roost, Marc M. Van Hulle

PII: S1053-8119(18)30294-5

DOI: 10.1016/j.neuroimage.2018.04.006

Reference: YNIMG 14853

To appear in: NeuroImage


Received Date: 13 June 2017

Revised Date: 22 January 2018

Accepted Date: 3 April 2018

Please cite this article as: Wittevrongel, B., Khachatryan, E., Hnazaee, M.F., Carrette, E., De Taeye, L., Meurs, A., Boon, P., Van Roost, D., Van Hulle, M.M., Representation of steady-state visual evoked potentials elicited by luminance flicker in human occipital cortex: An electrocorticography study, *NeuroImage* (2018), doi: 10.1016/j.neuroimage.2018.04.006.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Representation of steady-state visual evoked potentials elicited by luminance flicker in human occipital cortex: an electrocorticography study

Benjamin Wittevrongel^{*1}, Elvira Khachatryan¹, Mansoureh Fahimi Hnazaee¹, Evelien Carrette², Leen De Taeye², Alfred Meurs², Paul Boon², Dirk Van Roost³, and Marc M. Van Hulle¹

¹Laboratory for Neuro- and Psychophysiology, Department of Neuroscience, KU Leuven, Leuven, Belgium ²Laboratory of Clinical and Experimental Neurophysiology, Neurology Department, Ghent University Hospital, Ghent, Belgium ³Department of Neurosurgery, Ghent University Hospital, Ghent, Belgium

Abstract

Despite the widespread use of steady-state visual evoked potentials (SSVEPs) elicited by luminance flicker in clinical and research settings, their spatial and temporal representation in the occipital cortex largely remain elusive. We performed intracranial-EEG recordings in response to targets flickering at frequencies from 11 to 15 Hz using a subdural electrode grid covering the entire right occipital cortex of a human subject, and we were able to consistently locate the gazed stimulus frequency at the posterior side of the primary visual cortex (V1). Peripheral flickering, undetectable in scalp-EEG, elicited activations in the interhemispheric fissure at locations consistent with retinotopic maps. Both foveal and peripheral activations spatially coincided with activations in the high gamma band. We detected localized alpha synchronization at the lateral edge of V2 during stimulation and transient post-stimulation theta band activations at the posterior part of the occipital cortex. Scalp-EEG exhibited only a minor occipital post-stimulation theta activation, but a strong transient frontal activation.

keywords

SSVEP, ECoG, alpha, theta, gamma, fundamental, harmonic

^{*}corresponding author: benjamin.wittevrongel@kuleuven.be address: O&N2, Herestraat 49 bus 1021, 3000 Leuven, Belgium

دريافت فورى 🛶 متن كامل مقاله

- امکان دانلود نسخه تمام متن مقالات انگلیسی
 امکان دانلود نسخه ترجمه شده مقالات
 پذیرش سفارش ترجمه تخصصی
 امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
 امکان دانلود رایگان ۲ صفحه اول هر مقاله
 امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
 دانلود فوری مقاله پس از پرداخت آنلاین
 پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات
- ISIArticles مرجع مقالات تخصصی ایران