

Quality of life, functional status and adhesiolysis during elective abdominal surgery

Chema Strik, Martijn W.J. Stommel, Jeroen C. Hol, Harry van Goor, Richard P.G. ten Broek



PII: S0002-9610(17)30919-4

DOI: [10.1016/j.amjsurg.2017.08.001](https://doi.org/10.1016/j.amjsurg.2017.08.001)

Reference: AJS 12467

To appear in: *The American Journal of Surgery*

Received Date: 6 June 2017

Revised Date: 28 July 2017

Accepted Date: 7 August 2017

Please cite this article as: Strik C, Stommel MWJ, Hol JC, van Goor H, ten Broek RPG, Quality of life, functional status and adhesiolysis during elective abdominal surgery, *The American Journal of Surgery* (2017), doi: 10.1016/j.amjsurg.2017.08.001.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Abstract**

*Background:* Adhesiolysis during abdominal surgery can cause iatrogenic organ injury, increased operative time and a more complicated convalescence. We assessed the impact of adhesiolysis and adhesiolysis-related complications on quality of life and functional status following elective abdominal surgery

*Methods:* Prospective cohort study, comparing patients requiring and not requiring adhesiolysis during an elective laparotomy or laparoscopy using the SF-36 and DASl questionnaire scores.

*Results:* 518 patients were included. Pre- and postoperative quality of life did not significantly differ between both groups. Patients with adhesiolysis had a significantly lower pre- and postoperative functional status ( $p < 0.01$ ). Higher age, concomitant pulmonary disease, postoperative complications, readmissions and chronic abdominal pain 6 months after surgery were all associated with a significant and independent decline in quality of life and functional status six months after surgery.

*Conclusion and relevance:* Adhesiolysis in itself does not affect functional status and quality of life six months after surgery. Postoperative complications, readmissions and chronic abdominal pain are associated with a lower health status.

متن کامل مقاله

دریافت فوری ←

**ISI**Articles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات