


Author's Accepted Manuscript

Single dose of mirtazapine modulates whole-brain functional connectivity during emotional narrative processing

Emma Komulainen, Enrico Glerean, Katarina Meskanen, Roope Heikkilä, Lauri Nummenmaa, Tuukka T. Raij, Jari Lahti, Pekka Jylhä, Tarja Melartin, Erkki Isometsä, Jesper Ekelund



PII: S0925-4927(16)30336-5
DOI: <http://dx.doi.org/10.1016/j.psychresns.2017.03.009>
Reference: PSYN10663

To appear in: *Psychiatry Research: Neuroimaging*

Received date: 13 November 2016
Revised date: 17 February 2017
Accepted date: 20 March 2017

Cite this article as: Emma Komulainen, Enrico Glerean, Katarina Meskanen, Roope Heikkilä, Lauri Nummenmaa, Tuukka T. Raij, Jari Lahti, Pekka Jylhä, Tarja Melartin, Erkki Isometsä and Jesper Ekelund, Single dose of mirtazapine modulates whole-brain functional connectivity during emotional narrative processing, *Psychiatry Research: Neuroimaging* <http://dx.doi.org/10.1016/j.psychresns.2017.03.009>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and a review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Single dose of mirtazapine modulates whole-brain functional connectivity during emotional narrative processing

Emma Komulainen^a, Enrico Glerean^b, Katarina Meskanen^a, Roope Heikkilä^a, Lauri Nummenmaa^{b,c}, Tuukka T. Raij^{a,b,d}, Jari Lahti^{e,f,g}, Pekka Jylhä^{a,h}, Tarja Melartin^a, Erkki Isometsä^{a,h}, Jesper Ekelund^{a,i}.

^aUniversity of Helsinki and Helsinki University Hospital, Psychiatry, Helsinki, Finland

^bAalto University, School of Science, Department of Neuroscience and Biomedical Engineering, Espoo, Finland

^cTurku PET Centre and Department of Psychology, University of Turku, Finland

^dAalto NeuroImaging, Aalto University, Espoo, Finland

^eUniversity of Helsinki, Institute of Behavioral Sciences, Helsinki, Finland

^fFolkhälsan Research Center, Helsinki, Finland

^gHelsinki collegium of Advanced Studies, University of Helsinki, Finland

^hNational Institute of Health and Welfare, Department of Mental Health and Substance Abuse Services, Helsinki, Finland

ⁱVaasa Hospital District, Department of Psychiatry, Vaasa, Finland

Corresponding author:

Emma Komulainen

Department of Psychiatry, University of Helsinki, P.O. Box 590, FI-00029 HUS, Finland

E-mail: emma.komulainen@helsinki.fi

Phone: +358505203573

Fax number: +358947163735

Abstract

The link between neurotransmitter-level effects of antidepressants and their clinical effect remain poorly understood. A single dose of mirtazapine decreases limbic responses to fearful faces in

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات