

Accepted Manuscript

Vortical structures for nanomagnetic memory induced by dipole-dipole interaction in monolayer disks

Zhaosen Liu, Orion Ciftja, Xichao Zhang, Yan Zhou, Hou Ian

PII: S0749-6036(18)30152-6

DOI: [10.1016/j.spmi.2018.03.058](https://doi.org/10.1016/j.spmi.2018.03.058)

Reference: YSPMI 5588


To appear in: *Superlattices and Microstructures*

Received Date: 23 January 2018

Accepted Date: 22 March 2018

Please cite this article as: Z. Liu, O. Ciftja, X. Zhang, Y. Zhou, H. Ian, Vortical structures for nanomagnetic memory induced by dipole-dipole interaction in monolayer disks, *Superlattices and Microstructures* (2018), doi: 10.1016/j.spmi.2018.03.058.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Vortical Structures for Nanomagnetic Memory Induced by Dipole-Dipole Interaction in Monolayer Disks

Zhaosen Liu^{a1}, Orion Ciftja^b, Xichao Zhang^c, Yan Zhou^c, Hou Ian^{d,e2}

^a*Department of Applied Physics, Nanjing University of Information Science and Technology, Nanjing 210044, China*

^b*Department of Physics, Prairie View A&M University, Prairie View, TX 77446, USA*

^c*School of Science and Engineering, The Chinese University of Hong Kong, Shenzhen 518172, China*

^d*Institute of Applied Physics and Materials Engineering, University of Macau, Macau, China*

^e*UM Research Institute, Zhuhai, Guangdong, China*

Abstract

It is well known that magnetic domains in nanodisks can be used as storage units for computer memory. Using two quantum simulation approaches, we show here that spin vortices on magnetic monolayer nanodisks, which are chirality-free, can be induced by dipole-dipole interaction (DDI) on the disk-plane. When DDI is sufficiently strong, vortical and anti-vortical multi-domain textures can be generated simultaneously. Especially, a spin vortex can be easily created and deleted through either external magnetic or electrical signals, making them ideal to be used in nanomagnetic memory and logical devices. We demonstrate these properties in our simulations.

Keywords: Spin Vortex, Magnetic memory, Dipolar Interaction

1. Introduction

The long-range ferromagnetic order in a magnetic device tends to disappear when the energy due to thermal fluctuation becomes comparable to the anisotropic and exchange energy terms. This is a well known superparamagnetic limit imposed on the miniaturization of magnetic devices [1].

¹Email: liuzhsnj@yahoo.com

²Email: houian@umac.mo

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات