

Accepted Manuscript

Spin susceptibility as a test of unitary limit in disordered graphene systems

I. Grosu, T.-L. Biter

PII: S1386-9477(17)30822-6

DOI: [10.1016/j.physe.2017.12.013](https://doi.org/10.1016/j.physe.2017.12.013)

Reference: PHYSE 12984

To appear in: *Physica E: Low-dimensional Systems and Nanostructures*


Received Date: 6 June 2017

Revised Date: 7 November 2017

Accepted Date: 1 December 2017

Please cite this article as: I. Grosu, T.-L. Biter, Spin susceptibility as a test of unitary limit in disordered graphene systems, *Physica E: Low-dimensional Systems and Nanostructures* (2018), doi: 10.1016/j.physe.2017.12.013.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Spin susceptibility as a test of unitary limit in disordered graphene systems

I. Grosu and T.-L.Biter

Department of Physics, "Babes-Bolyai" University, 400084
Cluj-Napoca, Romania

We analyzed the static spin susceptibility of graphene systems in the presence of disorder and a small energy gap. We considered the case of strong scatterers (unitary limit). The temperature and impurity concentration effects were analyzed. The critical value of the impurity concentration was calculated. The behavior of the spin susceptibility for different values of the impurity concentration is discussed.

Key-words : Graphene, Disorder, Energy gap, Unitary limit, Critical disorder, Spin susceptibility.

I. Introduction

Graphene is a monolayer of carbon atoms packed in a two-dimensional lattice which was synthesized about a decade ago [1]. The band structure of graphene has linear dispersions near the two nonequivalent points named as K and K' , or Dirac points, of the Brillouin zone. Around the Dirac points, the kinetic energy for a bare electron is linear, $\varepsilon_{k,\lambda} = \lambda v_F |\vec{k}|$, where \vec{k} is the wave vector with respect to the

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات