

# Accepted Manuscript

Research papers

An improved bias correction method of daily rainfall data using a sliding window technique for climate change impact assessment

P.S. Smitha, B. Narasimhan, K.P. Sudheer, H. Annamalai

PII: S0022-1694(17)30764-3

DOI: <https://doi.org/10.1016/j.jhydrol.2017.11.010>

Reference: HYDROL 22368

To appear in: *Journal of Hydrology*


Received Date: 21 August 2017

Revised Date: 29 October 2017

Accepted Date: 7 November 2017

Please cite this article as: Smitha, P.S., Narasimhan, B., Sudheer, K.P., Annamalai, H., An improved bias correction method of daily rainfall data using a sliding window technique for climate change impact assessment, *Journal of Hydrology* (2017), doi: <https://doi.org/10.1016/j.jhydrol.2017.11.010>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



**An improved bias correction method of daily rainfall data using a sliding window technique for climate change impact assessment**

P. S. Smitha<sup>1</sup>, B. Narasimhan<sup>1</sup>, K. P. Sudheer<sup>1</sup> and H. Annamalai<sup>2</sup>

<sup>1</sup>Department of Civil Engineering, Indian Institute of Technology Madras, Chennai – 600036, India

<sup>2</sup>International Pacific Research Center and Department of Oceanography, University of Hawaii, USA.

**Abstract**

Regional climate models (RCMs) are used to downscale the coarse resolution General Circulation Model (GCM) outputs to a finer resolution for hydrological impact studies. However, RCM outputs often deviate from the observed climatological data, and therefore need bias correction before they are used for hydrological simulations. While there are a number of methods for bias correction, most of them use monthly statistics to derive correction factors, which may cause errors in the rainfall magnitude when applied on a daily scale. This study proposes a sliding window based daily correction factor derivations that help build reliable daily rainfall data from climate models. The procedure is applied to five existing bias correction methods, and is tested on six watersheds in different climatic zones of India for assessing the effectiveness of the corrected rainfall and the consequent hydrological simulations. The bias correction was performed on rainfall data downscaled using Conformal Cubic Atmospheric Model (CCAM) to  $0.5^\circ \times 0.5^\circ$  from two different CMIP5 models (CNRM-CM5.0, GFDL-CM3.0). The India Meteorological Department (IMD) gridded ( $0.25^\circ \times 0.25^\circ$ ) observed rainfall data was considered to test the effectiveness of the proposed bias correction method. The

متن کامل مقاله

دریافت فوری ←

**ISI**Articles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات