

Accepted Manuscript

Ferric ion pair mediated biomass redox flow fuel cell and related chemical reaction kinetics study

Xihong Zu, Lele Sun, Jian Gong, Xiaochun Liu, Yangxi Liu, Xu Du, Wei Liu, Lifen Chen, Guobin Yi, Weigang Zhang, Wenjing Lin, Weizhao Li, Yulin Deng

PII: S1385-8947(18)30548-5
DOI: <https://doi.org/10.1016/j.cej.2018.03.190>
Reference: CEJ 18797

To appear in: *Chemical Engineering Journal*

Received Date: 9 January 2018
Revised Date: 30 March 2018
Accepted Date: 31 March 2018

Please cite this article as: X. Zu, L. Sun, J. Gong, X. Liu, Y. Liu, X. Du, W. Liu, L. Chen, G. Yi, W. Zhang, W. Lin, W. Li, Y. Deng, Ferric ion pair mediated biomass redox flow fuel cell and related chemical reaction kinetics study, *Chemical Engineering Journal* (2018), doi: <https://doi.org/10.1016/j.cej.2018.03.190>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Ferric ion pair mediated biomass redox flow fuel cell and related chemical reaction kinetics study

Xihong Zu^a, Lele Sun^a, Jian Gong^b, Xiaochun Liu^a, Yangxi Liu^a, Xu Du^b, Wei Liu^b, Lifan Chen^a, Guobin Yi^{a,*}, Weigang Zhang^a, Wenjing Lin^a, Weizhao Li^a, Yulin Deng^{b,*}

^aSchool of Chemical Engineering and Light Industry, Guangdong University of Technology, Guangzhou, Guangdong, 510006, P.R. China

^bSchool of Chemical & Biomolecular Engineering, Georgia Institute of Technology, 500 10th Street N.W., Atlanta, GA 30332-0620, USA

Corresponding author: yulin.deng@ipst.gatech.edu; ygb702@163.com.

Abstract: Here, a novel and low cost redox flow fuel cell that directly converts raw biomass to electricity at low temperature without utilizing any noble metal catalyst is reported. In this cell, Fe³⁺ ion, which has strong oxidizing power and is chemical stable, was directly utilized as an oxidant for biomass in anolyte. The reduced Fe²⁺ acted as charge carriers, transferring the electrons to the anode of the cell and being converted back to Fe³⁺ simultaneously. The degradation chemistry of the biomass from the oxidation reaction and the apparent oxidation kinetics were investigated using glucose as the biomass model compound. The results showed that glucose was decomposed to small organic molecules and even CO₂, and the reaction rate was significantly affected by temperature and content of reactant. The apparent activation energy of the redox reaction is 127.85 kJ·mol⁻¹ based on the linear relation equation at the initial stage of the reaction. And the preference conditions of the redox reaction are 100~110 °C, 0.2~0.5 mol·L⁻¹ glucose and the mol ratio of

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات