



Contents lists available at ScienceDirect

Environmental Pollution

journal homepage: www.elsevier.com/locate/envpol

Temporal trends of infant and birth outcomes in Minamata after severe methylmercury exposure[☆]



Takashi Yorifuji^{a,*}, Saori Kashima^b, Made Ayu Hitapretiwu Suryadhi^a,
Kawuli Abudureyimu^a

^a Department of Human Ecology, Graduate School of Environmental and Life Science, Okayama University, 3-1-1 Tsushima-naka, Kita-ku, Okayama 700-8558, Japan

^b Department of Public Health and Health Policy, Graduate School of Biomedical and Health Sciences, Hiroshima University, 1-2-3 Kasumi, Minami-ku, Hiroshima 734-0037, Japan

ARTICLE INFO

Article history:

Received 23 June 2017

Received in revised form

13 September 2017

Accepted 19 September 2017

Available online 28 September 2017

Keywords:

Birth rate

Environmental pollution

Food contamination

Infant mortality

Methylmercury poisoning

Minamata disease

Sex ratio

Stillbirth

ABSTRACT

Severe methylmercury poisoning occurred in Minamata and neighboring communities in Japan during the 1950s and 1960s, causing what is known as Minamata disease. Although an increase in stillbirths and a reduced male proportion at birth (i.e., reduced sex ratio) have been reported, no studies have evaluated the impact of exposure on an entire set of infant and birth outcomes. We therefore evaluated the temporal trends of these outcomes in the Minamata area from 1950 to 1974. We focused on the spontaneous/artificial stillbirth rate, crude fertility rate, male proportion at birth, male proportion among stillbirths, and infant mortality. We obtained the number of stillbirths, live births, and infant deaths in Minamata City and Kumamoto Prefecture (as a reference) from 1950 to 1974. After plotting annual figures for each outcome, we divided the study period into five intervals and compared them between Minamata City and Kumamoto Prefecture using the chi-squared test. We observed a slightly increased spontaneous stillbirth rate and decreased artificial stillbirth rate in Minamata City, followed by a reduced crude fertility rate. The crude fertility rates in Minamata City during the period 1955–1965 were significantly lower compared with those in Kumamoto Prefecture ($p < 0.001$). An increase in the male proportion among stillbirths was observed, corresponding to a reduction in the proportion of males at birth in the late 1950s. The impact on infant mortality was equivocal. These descriptive analyses demonstrate a severe regional impact of methylmercury exposure on a series of birth outcomes in the Minamata area.

© 2017 Elsevier Ltd. All rights reserved.

1. Introduction

Severe methylmercury poisoning through consumption of contaminated seafood occurred in Minamata and neighboring communities in Japan during the 1950s and 1960s; the incident caused what is known as Minamata disease (Harada, 1995; Yorifuji et al., 2013a). The first patient was officially identified in May 1956, and many other patients were subsequently identified in the exposed areas. Affected individuals manifest neurological signs including paresthesia, ataxia, dysarthria, constriction of the visual

field, and hearing difficulties. Moreover, a considerable number of children who were exposed to methylmercury *in utero* were born in exposed areas during the exposure period with conditions resembling cerebral palsy, known as congenital Minamata disease (Harada, 1978).

Aside from congenital Minamata disease, exposure to methylmercury *in utero* in the contaminated areas affected several birth outcomes. For example, Itai et al. observed an increased proportion of abnormal pregnancies (spontaneous abortion and stillbirths) in a contaminated area of Minamata City compared with a control area (in a different prefecture) during the exposure period (Itai et al., 2004). Moreover, reduced sex ratio at birth (i.e., reduced male proportion) was observed in Minamata City and neighboring communities during the period (Doi et al., 1985; Sakamoto et al., 2001; Yorifuji and Kashima, 2016), which is considered to be owing to a greater proportion of male stillbirths (Sakamoto et al.,

[☆] This paper has been recommended for acceptance by Prof. W. Wen-Xiong.

* Corresponding author. Department of Human Ecology, Graduate School of Environmental and Life Science, Okayama University, 3-1-1 Tsushima-naka, Kita-ku, Okayama 700-8530, Japan.

E-mail address: yorichan@md.okayama-u.ac.jp (T. Yorifuji).




Fig. 1. Map of study area.

Table 1
Demographic characteristics of residents in Minamata City and Kumamoto Prefecture in 1960.

| | Exposure area | Reference |
|--|-------------------------------|--|
| | Minamata City (n = 48,342) | Kumamoto prefecture (n = 1,856,192) |
| Age category, n (%) | | |
| 0–14 years old | 17,339 (35.9) | 629,548 (33.9) |
| 15–59 years old | 26,833 (55.5) | 1,033,097 (55.7) |
| 60 years old and over | 4170 (8.6) | 193,547 (10.4) |
| Women, n (%) | 25,270 (52.3) | 969,154 (52.2) |
| Women aged 15–44 years among all women, n (%) | 11,163 (44.2) | 421,221 (43.5) |
| Educational status (among residents aged 15 years old and over) | | |
| Number of residents aged 15 years old and over, n | 31,003 | 1,226,644 |
| Residents who completed university, n (%) | 418 (1.3) | 14,724 (1.2) |
| Residents who completed high school or college, n (%) | 2772 (8.9) | 107,493 (8.8) |
| Industry classification (among workers aged 15 years old and over) | | |
| Number of workers aged 15 years old and over, n | 19,944 | 814,544 |
| Primary (Farmer, fisherman, etc.), n (%) | 6283 (31.5) | 420,310 (51.6) |
| Secondary (Factory worker, etc.), n (%) | 6510 (32.6) | 133,923 (16.4) |
| Tertiary (Services, etc.), n (%) | 7146 (35.8) | 260,235 (31.9) |
| Unknown, n | 5 | 76 |

Source: National census (1960).

2001). However, no studies thus far have evaluated the impact of exposure on an entire set of infant and birth outcomes (e.g., crude fertility rate or infant mortality rate) in the contaminated area.

We therefore conducted a descriptive study to evaluate

temporal trends of infant and birth outcomes in the Minamata area from 1950 to 1974 and assessed how regional methylmercury exposure affected these outcomes.

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات