



Featured Article

Progress toward standardized diagnosis of vascular cognitive impairment: Guidelines from the Vascular Impairment of Cognition Classification Consensus Study

Olivia A. Skrobot^a, Sandra E. Black^b, Christopher Chen^c, Charles DeCarli^d, Timo Erkinjuntti^e, Gary A. Ford^f, Rajesh N. Kalaria^g, John O'Brien^h, Leonardo Pantoniⁱ, Florence Pasquier^j, Gustavo C. Roman^k, Anders Wallin^l, Perminder Sachdev^{m,n}, Ingmar Skoog^o, VICCCS group¹, Yoav Ben-Shlomo^p, Anthony P. Passmore^q, Seth Love^a, Patrick G. Kehoe^{a,*}

^aDementia Research Group, School of Clinical Sciences, Faculty of Health Sciences, University of Bristol, Level 1, Learning & Research, Southmead Hospital, Bristol, UK

^bSunnybrook Research Institute, University of Toronto, Canada

^cMemory Aging & Cognition Centre, Department of Pharmacology, National University of Singapore, Singapore

^dAlzheimer's Disease Center and Imaging of Dementia and Aging (IDEA) Laboratory, Department of Neurology and Center for Neuroscience, University of California at Davis, 4860 Y Street, Suite 3700, Sacramento, CA 95817, USA

^eClinical Neurosciences, Neurology, University of Helsinki and Helsinki University Hospital, Finland, POB 300, FIN-00290, HUS, Finland

^fDivision of Medical Sciences, Oxford University, Magdalen Centre North, Oxford Science Park, OX4 4GA, UK

^gInstitute of Neuroscience, Campus for Ageing & Vitality Newcastle upon Tyne, NIHR Biomedical Research Building, NE4 5PL, UK

^hDepartment of Psychiatry, University of Cambridge School of Clinical Medicine, Cambridge Biomedical Campus Box 189, Level E4, Cambridge, CB2 0SP UK

ⁱNEUROFARBA Department, University of Florence, Florence, Italy

^jUniversity of Lille, Inserm U1171, Degenerative and vascular disorders, CHU, Distal, F-59000, Lille, France

^kMethodist Neurological Institute, 6560 Fannin Street, Suite 802, Houston, TX 77030, USA

^lInstitute of Neuroscience and Physiology at Sahlgrenska Academy, University of Gothenburg, Memory Clinic at Department of Neuropsychiatry, Sahlgrenska University Hospital, Wallingatan 6, SE-431 41 Mölndal, Sweden

^mSchool of Psychiatry, University of New South Wales, Sydney, Australia

ⁿCHeBA (Centre for Healthy Brain Ageing), Neuropsychiatric Institute, Prince of Wales Hospital, Randwick NSW 2031, Australia

^oCenter for Health and Ageing (AGECAP), Institute of Neuroscience and Physiology, Sahlgrenska Academy, University of Gothenburg, Sweden

^pSchool of Social and Community Medicine, University of Bristol, Canynge Hall, 39 Whatley Road, Bristol, BS8 2PS, UK

^qInstitute of Clinical Sciences, Block B, Queens University Belfast, Royal Victoria Hospital, Belfast, BT12 6BA

Abstract

Introduction: Progress in understanding and management of vascular cognitive impairment (VCI) has been hampered by lack of consensus on diagnosis, reflecting the use of multiple different assessment protocols. A large multinational group of clinicians and researchers participated in a two-phase Vascular Impairment of Cognition Classification Consensus Study (VICCCS) to agree on principles (VICCCS-1) and protocols (VICCCS-2) for diagnosis of VCI. We present VICCCS-2.

Methods: We used VICCCS-1 principles and published diagnostic guidelines as points of reference for an online Delphi survey aimed at achieving consensus on clinical diagnosis of VCI.

Results: Six survey rounds comprising 65–79 participants agreed guidelines for diagnosis of VICCCS-revised mild and major forms of VCI and endorsed the National Institute of Neurological Disorders–Canadian Stroke Network neuropsychological assessment protocols and recommendations for imaging.

¹Members are listed at the end of the article.

E-mail address: patrick.kehoe@bristol.ac.uk

*Corresponding author. Tel.: 0117 414 7821.

Discussion: The VICCCS-2 suggests standardized use of the National Institute of Neurological Disorders–Canadian Stroke Network recommendations on neuropsychological and imaging assessment for diagnosis of VCI so as to promote research collaboration.
© 2017 the Alzheimer's Association. Published by Elsevier Inc. All rights reserved.

Keywords: Vascular cognitive impairment; Vascular dementia; Guidelines; Criteria; consensus; Delphi

1. Introduction

Since Hachinski et al [1] proposed the term multi-infarct dementia to describe dementia complicating ischemic vascular disease, numerous other descriptors have been used to encompass the heterogeneous clinical and etiological spectrum of cognitive impairment due to cerebrovascular disease (CVD). These include vascular dementia (VaD), vascular cognitive impairment (VCI), subcortical (ischemic) vascular dementia, and vascular cognitive disorders (VCDs), variably diagnosed according to multiple different guidelines or protocols [2–14], some agreed by national institutions or research networks, for example, Alzheimer's Disease Diagnostic and Treatment Centers [11], International Statistical Classification of Diseases, 10th revision [15], the National Institute of Neurological Disorders and Stroke (NINDS)–Association Internationale pour la Recherche et l'Enseignement en Neurosciences [16], and Diagnostic and Statistical Manual of Mental Disorders, fourth and fifth editions (DSM-4 and DSM-5; [17,18]).

Studies comparing some of these protocols have shown they are not readily interchangeable [19–21]. After the commencement of the Vascular Impairment of Cognition Classification Consensus Study phase 1 (VICCCS-1), the American Heart Association/American Stroke Association (AHA/ASA) published a statement on vascular contributions to cognitive impairment and dementia [22]. This supported the use of assessment protocols previously published by NINDS–Canadian Stroke Network (CSN) [13]. There have been other recent contributions to this field from the International Society of Vascular Behavioral and Cognitive Disorders (VASCOD) [23] and the DSM-5 [18]. The level of take up of these recent guidelines is still unclear. Only those published during VICCCS-1, before commencement of VICCCS-2, could be included for consideration in the present study [22,24].

The aim of VICCCS was to achieve broad international consensus on diagnosis of VCI, through participation of a large pool of international researchers and clinicians in an iterative survey using the Delphi approach. After two initial survey rounds, the study was separated into two phases: VICCCS-1, addressing key concepts in our understanding and terminology of cognitive impairment resulting from CVD [25], and VICCCS-2, focusing on the formulation of practical guidelines for diagnosis.

The VICCCS-1 achieved broad consensus on concepts of VCI. It supported the use of “mild” and “major” subdivisions

of the severity of impairment, aligning with the revised terminology in the DSM-5. VICCCS-1 participants concluded that attempts to separate mild VCI into further subtypes according to affected cognitive domains were at present premature but agreed that this should be an area of future research. VICCCS-1 agreed (Fig. 1, reproduced from [25]) that the major forms of VCI (VaD) should be classified into four main subtypes: (i) post-stroke dementia (PSD); (ii) subcortical ischemic vascular dementia (SIVaD); (iii) multi-infarct (cortical) dementia (MID); and (iv) mixed dementias (further subdivided according to additional neurodegenerative pathologies). Framed by these concepts, VICCCS-2 used the same Delphi methodology to agree diagnostic guidelines on determination of severity of VCI and discrimination of subtypes.

2. Methods

Participants in VICCCS-1 [25] were invited to participate in VICCCS-2 (Supplementary Figure 1). Although 149 initially agreed to participate, only approximately half were active and committed respondents in three or more rounds, with low attrition and little variation in participation throughout the six rounds (65–79 participants in each round, a mean of 72). Of the active participants, 63%–75% of participants (mean 68%) were clinicians with direct involvement in clinical assessment or health service patient care. The remainder were nonclinical (i.e., supporting clinical work technically or otherwise, but not involved in clinical decision-making, or predominantly involved in research). Individual round representation is provided in Supplementary Table 1.

2.1. Data collection

We used the Delphi method, an iterative, multistaged series of structured questionnaires with feedback of anonymized responses and progressive refinement of questions to reach consensus [26]. The process was co-ordinated by a nonparticipating researcher (O.A.S). Anonymization of responses facilitated free expression of opinion throughout the study. Feedback of summary responses after each round informed subsequent questions and allowed unbiased evolution of group judgment. A threshold of two-thirds agreement was chosen to signify consensus [27] for issues refined iteratively through multiple rounds, as in VICCCS-1 [25]. For issues where this threshold was not reached, we present

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات