Appl. Comput. Harmon. Anal. ••• (••••) •••-••

ELSEVIER

Contents lists available at ScienceDirect

# Applied and Computational Harmonic Analysis

Applied and Computational Harmonic Analysis

www.elsevier.com/locate/acha

## On the number of harmonic frames

Simon Marshall<sup>a</sup>, Shayne Waldron<sup>b,\*</sup>

<sup>a</sup> Department of Mathematics, University of Wisconsin-Madison, 480 Lincoln Drive, Madison, WI 53706, IISA

b Department of Mathematics, University of Auckland, Private Bag 92019, Auckland, New Zealand

#### ARTICLE INFO

### Article history: Received 21 November 2016 Received in revised form 17 October 2017 Accepted 6 February 2018

Accepted 6 February 2018 Available online xxxx Communicated by Zuowei Shen

MSC: primary 42C15, 94A12 secondary 11Z05, 20C15, 94A15, 20F28

Keywords:
Finite tight frames
Harmonic frames
Finite abelian groups
Character theory
Pontryagin duality
Mordell-Lang conjecture
Projective unitary equivalence

#### ABSTRACT

There is a finite number  $h_{n,d}$  of tight frames of n distinct vectors for  $\mathbb{C}^d$  which are the orbit of a vector under a unitary action of the cyclic group  $\mathbb{Z}_n$ . These cyclic harmonic frames (or geometrically uniform tight frames) are used in signal analysis and quantum information theory, and provide many tight frames of particular interest. Here we investigate the conjecture that  $h_{n,d}$  grows like  $n^{d-1}$ . By using a result of Laurent which describes the set of solutions of algebraic equations in roots of unity, we prove the asymptotic estimate

$$h_{n,d} \approx \frac{n^d}{\varphi(n)} \ge n^{d-1}, \qquad n \to \infty.$$

By using a group theoretic approach, we also give some exact formulas for  $h_{n,d}$ , and estimate the number of cyclic harmonic frames up to projective unitary equivalence. © 2018 Elsevier Inc. All rights reserved.

#### 1. Introduction

Tight frames of n vectors for  $\mathbb{C}^d$  have numerous applications (see the surveys [4], [5]). These include signal transmission with erasures [10], [14], [2] and quantum information theory [16], [20].

Many tight frames of practical and theoretical interest are G-frames (the orbit of a unitary action of a group G) [22]. Most notable are the harmonic frames (G is abelian) and SICs, i.e.,  $d^2$  equiangular lines in  $\mathbb{C}^d$  (for a projective action of the abelian group  $\mathbb{Z}^2_d$ ). The main result of this paper is a precise statement about how numerous the harmonic frames of n vectors for  $\mathbb{C}^d$  are (Theorem 3.1). By way of comparison, SICs are known to exist only for certain values of d, and there is strong evidence for Zauner's conjecture that they exist for all values of d (see [20], [25]).

E-mail address: waldron@math.auckland.ac.nz (S. Waldron).

https://doi.org/10.1016/j.acha.2018.02.004

1063-5203/© 2018 Elsevier Inc. All rights reserved.

<sup>\*</sup> Corresponding author.

2

We now provide some background on harmonic frames, and then detail our approach (precise definitions are given in §2). What we will call a *cyclic harmonic frame* for  $\mathbb{C}^d$  was first introduced as a  $d \times n$  submatrix  $[v_1, \ldots, v_n]$  of the Fourier matrix (character table for  $\mathbb{Z}_n$ )

$$\begin{bmatrix} 1 & 1 & 1 & \cdots & 1 \\ 1 & \omega & \omega^{2} & \cdots & \omega^{n-1} \\ 1 & \omega^{2} & \omega^{4} & \cdots & \omega^{2(n-1)} \\ \vdots & \vdots & \vdots & & \vdots \\ 1 & \omega^{n-1} & \omega^{2(n-1)} & \cdots & \omega^{(n-1)(n-1)} \end{bmatrix}, \qquad \omega := e^{\frac{2\pi i}{n}}, \tag{1.1}$$

obtained by selecting d of the rows (characters of  $\mathbb{Z}_n$ ). See [11], [13], [3] (who use the term harmonic frame for when the first d rows are taken), [6] (who use the term a Fourier ensemble), and [19] (who show a matrix with its columns given by a random cyclic harmonic frame is a RIP (restricted isometry property) matrix with high probability). These tight frames can be viewed as  $\mathbb{Z}_n$ -frames (called geometrically uniform frames in [9]). This construction generalises, with  $\mathbb{Z}_n$  replaced by an abelian group G of order n [21], to give what we call a harmonic frame (it is cyclic if G can be taken to be  $\mathbb{Z}_n$ ). It follows from the character table construction (and the fact there are a finite number of abelian groups of order n) that there is a finite number of harmonic frames of n vectors for  $\mathbb{C}^d$ .

A computer study [24] of the harmonic frames of n vectors for  $\mathbb{C}^d$  suggested the following behaviour:

- The number of harmonic frames (up to unitary equivalence) grows like  $n^{d-1}$ , and it is influenced by the prime factors of n.
- The majority of harmonic frames are cyclic.

In this paper, we show that for fixed d the number  $h_{n,d}$  of cyclic harmonic frames grows like

$$h_{n,d} \approx \frac{n^d}{\varphi(n)} \ge n^{d-1}, \quad n \to \infty.$$

The key points of our argument are

- Cyclic harmonic frames correspond to d-element subsets  $J \subset \mathbb{Z}_n$ .
- When cyclic harmonic frames given by  $J, K \subset \mathbb{Z}_n$  are unitarily equivalent, usually  $K = \sigma J$  for some automorphism. When this is not the case, we say they are *exceptional*.
- The automorphisms of  $\mathbb{Z}_n$  are easy to describe (as the units  $\mathbb{Z}_n^*$ ).
- A pair of unitarily equivalent cyclic harmonic frames determines a torsion point on the (2d)-torus  $\mathbb{T}^{2d}$ .
- By using results about the torsion point solutions of algebraic equations, we show that the number of exceptional harmonic frames grows slower than the number which aren't.
- The nonexceptional cyclic harmonic frames are counted by Burnside enumeration.

We carry out this argument in §4–§5. We give examples and some numerical data in §6. In §7 we show that there are no exceptional equivalences when n is prime, and together with Burnside enumeration this allows us to give an exact formula for  $h_{n,d}$  in this case, which we break down into lifted and unlifted, and into real and complex harmonic frames.

In the final section, we use our techniques to investigate the number  $p_{n,d}$  of harmonic frames of n vectors for  $\mathbb{C}^d$  up to projective unitary equivalence. For  $d \geq 4$ , this gives the lower estimate

$$p_{n,d} \approx \frac{n^{d-1}}{\varphi(n)} \ge n^{d-2}, \quad n \to \infty.$$

# دريافت فورى ب متن كامل مقاله

# ISIArticles مرجع مقالات تخصصی ایران

- ✔ امكان دانلود نسخه تمام متن مقالات انگليسي
  - ✓ امكان دانلود نسخه ترجمه شده مقالات
    - ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
  - ✓ امكان دانلود رايگان ۲ صفحه اول هر مقاله
  - ✔ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
    - ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات