



ELSEVIER

Contents lists available at ScienceDirect

Journal of Number Theory

www.elsevier.com/locate/jnt



The Roman harmonic numbers revisited

J. Sesma

Departamento de Física Teórica, Facultad de Ciencias, 50009, Zaragoza, Spain

ARTICLE INFO

Article history:

Received 14 February 2017
Received in revised form 31 March 2017

Accepted 15 May 2017

Available online xxxx

Communicated by the Principal Editors

MSC:

11B75
11B73
11B68

Keywords:

Generalized harmonic numbers
Quantum distribution moments
Stirling numbers
Generating relations
Sum rules
Pochhammer symbol

ABSTRACT

Two decades ago, Steven Roman, Daniel E. Loeb and Gian-Carlo Rota introduced a family of harmonic numbers in their study of harmonic logarithms. We propose to refer to those numbers as *Roman harmonic numbers*. With the purpose of revitalizing the study of these mathematical objects, we recall here their known properties and unveil additional ones. An integral representation, several generating relations, and a collection of sum rules involving those numbers are presented. It is also shown that higher derivatives of the Pochhammer and reciprocal Pochhammer symbols are easily expressed in terms of Roman harmonic numbers.

© 2017 Elsevier Inc. All rights reserved.

1. Introduction

In the course of a research on the so called epsilon expansions of Appell and Kampé de Fériet functions [11], we considered convenient to define what we called modified generalized harmonic numbers, $\hat{H}_n^{(k)}$, in the form

E-mail address: javier@unizar.es.

<http://dx.doi.org/10.1016/j.jnt.2017.05.009>

0022-314X/© 2017 Elsevier Inc. All rights reserved.

$$\hat{H}_0^{(k)} = \delta_{k,0}, \quad \hat{H}_n^{(k)} \equiv \sum_{j=1}^n (-1)^{j-1} \binom{n}{j} \frac{1}{j^k}, \quad n \geq 1. \quad (1)$$

Later on, we realized that these mathematical objects, with the name of harmonic numbers and denoted by $c_n^{(k)}$, had already been introduced, by Loeb and Rota [16] and by Roman [22,23], to give explicit expressions of the harmonic logarithms. On the other hand, the moments

$$\bar{d}^p = \int_0^\infty d\delta \delta^p \mathcal{P}(\delta) \quad p = 0, 1, 2, \dots, \quad (2)$$

of the quantum probability distribution

$$\mathcal{P}(\delta) = 2(D_\varepsilon - 1) e^{-2\delta} (1 - e^{-2\delta})^{D_\varepsilon - 2}, \quad D_\varepsilon = 2, 3, \dots, \quad (3)$$

discussed by Coffey [7] and used (for $p = 1$ and 2) to approximately calculate the decoherence factors for D_ε -dimensional quantum states, are closely related to the same harmonic numbers, as shown below. In spite of their interesting properties, the $c_n^{(k)}$ have not received, in our opinion, due attention by number theorists. Here we intend to arouse the interest in those harmonic numbers by recalling their known properties and presenting some others.

Several generalizations of harmonic numbers have been used by different authors [2–6,10,14,15,24]. To avoid misunderstandings, we adopt the name of *Roman harmonic numbers* for those introduced in Refs. [16,22], and [23]. One of the possible equivalent definitions of these numbers is given in Section 2, which recalls also their main properties, found by Roman [22,23] and by Loeb and Rota [16]. Section 3 shows, firstly, the connection of those numbers with known nested sums. Their integral representation allows to discover the relation between Roman harmonic numbers and Coffey's quantum distribution moments. Then, several generating relations, sum rules, and additional properties not mentioned before are presented. As an application, expressions are given of the derivatives of the Pochhammer and reciprocal Pochhammer symbols in terms of the $c_n^{(k)}$. To end, some pertinent comments are included in Section 4.

2. First definitions and known properties

Before recalling the Roman harmonic numbers, some auxiliary definitions, given in [16, 22], and [23], are necessary. We present them with the notation used in these references.

Definition 1. (Roman number.) The *Roman* n is defined to be

$$[n] = \begin{cases} n & \text{for } n \neq 0, \\ 1 & \text{for } n = 0. \end{cases} \quad (4)$$

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات