



Exponential stability of linear delayed differential systems



Leonid Berežansky^a, Josef Diblík^{b,*}, Zdeněk Svoboda^b, Zdeněk Šmarda^b

^a Department of Mathematics, Ben-Gurion University of the Negev, Beer-Sheva 84105, Israel

^b Brno University of Technology, Department of Mathematics, Faculty of Electrical Engineering and Communication, Brno, Czech Republic

ARTICLE INFO

MSC:
34K20
34K06
34K25

Keywords:

Linear delayed differential system
Exponential stability
Bohl–Perron theorem

ABSTRACT

Linear delayed differential systems

$$\dot{x}_i(t) = - \sum_{j=1}^m \sum_{k=1}^{r_{ij}} a_{ij}^k(t) x_j(h_{ij}^k(t)), \quad i = 1, \dots, m$$

are analyzed on a half-infinity interval $t \geq 0$. It is assumed that m and r_{ij} , $i, j = 1, \dots, m$ are natural numbers and the coefficients $a_{ij}^k : [0, \infty) \rightarrow \mathbb{R}$ and delays $h_{ij}^k : [0, \infty) \rightarrow \mathbb{R}$ are measurable functions. New explicit results on uniform exponential stability are derived including, as partial cases, recently published results.

© 2017 Elsevier Inc. All rights reserved.

1. Introduction

There are two main forms of linear systems of delay differential equations. The first one is a scalar form

$$\dot{x}_i(t) = - \sum_{j=1}^m \sum_{k=1}^{r_{ij}} a_{ij}^k(t) x_j(h_{ij}^k(t)), \quad i = 1, \dots, m, \quad (1)$$

where $t \geq t_0 \geq 0$, $t_0 \in \mathbb{R}$, m and r_{ij} , $i, j = 1, \dots, m$ are natural numbers, coefficients $a_{ij}^k : [0, \infty) \rightarrow \mathbb{R}$ and delays $h_{ij}^k : [0, \infty) \rightarrow \mathbb{R}$ are given functions, and $x_i : [-h, \infty) \rightarrow \mathbb{R}$, where

$$h := \inf_{t \geq 0} \min_{\substack{i, j=1, \dots, m \\ k=1, \dots, r_{ij}}} \{h_{ij}^k(t)\},$$

are unknown functions.

The second one is a vector form

$$\dot{X}(t) = - \sum_{k=1}^m A_k(t) X(h_k(t)), \quad (2)$$

where $t \geq t_0 \geq 0$, $t_0 \in \mathbb{R}$, m is a natural number, $A_k = (a_{ij}^k)_{i, j=1}^m : [0, \infty) \rightarrow \mathbb{R}^{2m}$ are matrix functions, $h_k : [0, \infty) \rightarrow \mathbb{R}$ are delay functions, and

$$X = (x_1, \dots, x_m)^T : [h^*, \infty) \rightarrow \mathbb{R}^m,$$

* Corresponding author.

E-mail addresses: brznsky@cs.bgu.ac.il (L. Berežansky), diblik@feec.vutbr.cz (J. Diblík), svobodaz@feec.vutbr.cz (Z. Svoboda), smarda@feec.vutbr.cz (Z. Šmarda).

where

$$h^* := \inf_{t \geq 0} \min_{i=1, \dots, m} \{h_i(t)\},$$

is an unknown vector function.

Papers on stability of linear differential delayed system can also be divided into two groups. The larger one considers vector form (2) (see for example papers [1–6] and monographs [7,8]) while the smaller one is devoted to scalar systems (1), we refer to papers [9–13] and to a monograph [14].

In the paper, we will study scalar systems (1). First, let us review some known stability results. In order to simplify this review, we consider non-autonomous systems and do not include the results specified for autonomous systems only (for such systems, we refer to interesting papers [11,13]). We also do not consider results obtained under the assumption

$$h_{ii}^k(t) \equiv t, \quad i = 1, \dots, m, \quad k = 1, \dots, r_{ii}.$$

In the sequel, we frequently use the concept of a non-singular M -matrix. For the sake of convenience, we recall this notion.

Definition 1 ([15]). An $m \times m$ matrix $G = (g_{ij})_{i,j=1}^m$ is called a non-singular M -matrix if $g_{ij} \leq 0, i, j = 1, \dots, m, i \neq j$ and one of the following equivalent conditions holds:

1. There exists a positive inverse matrix G^{-1} .
2. All the principal minors of matrix G are positive.

In [12], the authors consider a non-autonomous system, which is a partial case of system (1),

$$\dot{x}_i(t) = - \sum_{j=1}^m a_{ij}(t)x_j(h_{ij}(t)), \quad i = 1, \dots, m, \tag{3}$$

where $t \in [t_0, \infty), a_{ij}(t), h_{ij}(t)$ are continuous functions, $h_{ij}(t) \leq t$, and $h_{ij}(t)$ are monotone increasing functions such that $\lim_{t \rightarrow \infty} h_{ij}(t) = \infty, i, j = 1, \dots, m$.

Theorem 1. [12, Theorem 2.2] Assume that, for $t \geq t_0$, there exist non-negative numbers $b_{ij}, i, j = 1, \dots, m, i \neq j$ such that $|a_{ij}(t)| \leq b_{ij}a_{ii}(t), i, j = 1, \dots, m, i \neq j, a_{ii}(t) \geq 0$ and

$$\int_{h_{ii}(t)}^{\infty} a_{ii}(s)ds = \infty, \quad d_i = \limsup_{t \rightarrow \infty} \int_{h_{ii}(t)}^t a_{ii}(s)ds < 3/2, \quad i = 1, \dots, m.$$

Let, for the entries $\tilde{b}_{ij}, i, j = 1, \dots, m$ of an $m \times m$ matrix \tilde{B} ,

$$\tilde{b}_{ii} = 1, \quad i = 1, \dots, m$$

and, for $i \neq j, i, j = 1, \dots, m$,

$$\tilde{b}_{ij} = \begin{cases} -\left(\frac{2 + d_i^2}{2 - d_i^2}\right)b_{ij}, & \text{if } d_i < 1, \\ -\left(\frac{1 + 2d_i}{3 - 2d_i}\right)b_{ij}, & \text{if } d_i \geq 1. \end{cases}$$

If \tilde{B} is a non-singular M -matrix, then system (3) is asymptotically stable.

Theorem 1 is an object of attention since, for a scalar case it coincides with the well-known 3/2 stability condition (we refer, e.g., to [16–18]). Nevertheless, we point out the following restrictions in **Theorem 1**. The result is valid only for systems with continuous coefficients and delays and the proof of the theorem depends on this assumption. Moreover, any equation of system (3) includes only one diagonal term. It means that system (3) is a partial case of (1) for $r_{ij} = 1, i, j = 1, \dots, m$. Besides, delays in system (1) must be monotone increasing functions.

In monograph [14], the following result for the system

$$\dot{x}_i(t) = - \sum_{k=1}^m \sum_{j=1}^n a_{ij}^k(t)x_j(h_k(t)), \quad i = 1, \dots, n \tag{4}$$

is proved. In its formulation, we use the notation $a^+ = \max\{a, 0\}, a \in \mathbb{R}$ and, for a function f essentially bounded on $[t_0, \infty)$, we use the norm

$$\|f\|_{L^\infty} = \text{ess sup}_{t \geq t_0} \|f(t)\|.$$

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات